





SELECTING  
PLANTS  
FOR  
POLLINATORS



A REGIONAL GUIDE FOR FARMERS, LAND MANAGERS, AND GARDENERS IN THE



**CASCADE  
MIXED FOREST  
CONIFEROUS  
FOREST  
ALPINE MEADOW  
PROVINCE**

INCLUDING THE STATES OF:

OREGON  
AND  
WASHINGTON

**POLLINATOR  
PARTNERSHIP**

and  
**NAPPC**

---

# TABLE OF CONTENTS

---

<b>WHY SUPPORT POLLINATORS?</b>	<b>4</b>
<b>GETTING STARTED</b>	<b>5</b>
<b>CASCADE MIXED FOREST</b>	<b>6</b>
<b>MEET THE POLLINATORS</b>	<b>8</b>
<b>PLANT TRAITS</b>	<b>10</b>
<b>DEVELOPING PLANTINGS</b>	<b>12</b>
<b>FARMS</b>	<b>13</b>
<b>PUBLIC LANDS</b>	<b>14</b>
<b>HOME LANDSCAPES</b>	<b>15</b>
<b>BLOOM PERIODS</b>	<b>16</b>
<b>PLANTS THAT ATTRACT POLLINATORS</b>	<b>18</b>
<b>HABITAT HINTS</b>	<b>20</b>
<b>CHECKLIST</b>	<b>22</b>
<b>RESOURCES AND FEEDBACK</b>	<b>23</b>

This is one of several guides for different regions in the United States. We welcome your feedback to assist us in making the future guides useful. Please contact us at [feedback@pollinator.org](mailto:feedback@pollinator.org)

Cover bee photo by Diane Petit

# SELECTING PLANTS FOR POLLINATORS

## **A REGIONAL GUIDE FOR FARMERS, LAND MANAGERS, AND GARDENERS**

### IN THE ECOLOGICAL REGION OF THE **CASCADE MIXED FOREST CONIFEROUS FOREST ALPINE MEADOW PROVINCE**

INCLUDING THE STATES OF:

OREGON

AND

WASHINGTON

A NAPPC AND POLLINATOR PARTNERSHIP™ PUBLICATION

---

This guide was funded by the National Fish and Wildlife Foundation, the C.S. Fund, the Plant Conservation Alliance, the U.S. Forest Service, and the Bureau of Land Management with oversight by the Pollinator Partnership™ ([www.pollinator.org](http://www.pollinator.org)), in support of the North American Pollinator Protection Campaign (NAPPC—[www.nappc.org](http://www.nappc.org)).



## WHY SUPPORT POLLINATORS?

IN THEIR 1996 BOOK, *THE FORGOTTEN POLLINATORS*, Buchmann and Nabhan estimated that animal pollinators are needed for the reproduction of 90% of flowering plants and one third of human food crops. Each of us depends on these industrious pollinators in a practical way to provide us with the wide range of foods we eat. In addition, pollinators are part of the intricate web that supports the biological diversity in natural ecosystems that helps sustain our quality of life.

Abundant and healthy populations of pollinators can improve fruit set and quality, and increase fruit size. In farming situations this increases production per acre. In the wild, biodiversity increases and wildlife food sources increase.

Alfalfa, cucumbers, raspberries, apples, and strawberries are some of the crops raised in the Cascade Mixed Forest that rely on honey bees and native bees for pollination. Domestic honey bees pollinate approximately \$10 billion worth of crops in the U.S. each year.

Unfortunately, the numbers of both native pollinators and domesticated bee populations are declining. They are threatened by habitat loss, disease, and the excessive and inappropriate use of pesticides. The loss of commercial bees to Colony Collapse Disorder (CCD) has highlighted how severe the issues of proper hive management are to reduce stresses caused by disease, pesticide use, insufficient nutrition, and transportation practices. Currently, the pollination services that the commercial beekeeping industry provides are receiving much needed research and conservation resources. The efforts to understand the threats to commercial bees should help us understand other pollinators and their roles in the environment as well.

It is imperative that we take immediate steps to help pollinator populations thrive. The beauty of the situation is that by supporting pollinators' need for habitat, we support our own needs for food and support diversity in the natural world.

Thank you for taking time to consult this guide. By adding plants to your landscape that provide food and shelter for pollinators throughout their active seasons and by adopting pollinator friendly landscape practices, you can make a difference to both the pollinators and the people that rely on them.



**Laurie Davies Adams**  
Executive Director  
Pollinator Partnership

“**FARMING FEEDS  
THE WORLD, AND  
WE MUST REMEMBER  
THAT POLLINATORS  
ARE A CRITICAL  
LINK IN OUR FOOD  
SYSTEMS.**”

-- PAUL GROWALD,  
CO-FOUNDER,  
POLLINATOR PARTNERSHIP



THIS REGIONAL GUIDE IS just one in a series of plant selection tools designed to provide information on how individuals can influence pollinator populations through choices they make when they farm a plot of ground, manage large tracts of public land, or plant a garden. Each of us can have a positive impact by providing the essential habitat requirements for pollinators including food, water, shelter, and enough space to allow pollinators to raise their young.

Pollinators travel through the landscape without regard to property ownership or state boundaries. We've chosen to use R.G. Bailey's classification system to identify the geographic focus of this guide and to underscore the connections between climate and vegetation types that affect the diversity of pollinators in the environment.

Bailey's Ecoregions of the United States, developed by the United States Forest Service, is a system created as a management tool

and is used to predict responses to land management practices throughout large areas. This guide addresses pollinator-friendly land management practices in what is known as the Cascade Mixed Forest, Coniferous Forest, Alpine Meadow Province.


Portions of Oregon and Washington make up the 53,400 square miles of this province with elevations ranging from sea level to over 14,000 feet. The topography is primarily mountainous. The Cascade Province includes steep mountains bordered by a narrow coastal plain, and the interior Cascade Range features mountains of 8,000 to 9,000 feet, punctuated every 5-85 miles by a much higher volcano. Average annual temperatures are mild, ranging from 35° to 50°F.

This wet, green province is characterized by heavy rains in winter, with summer fog, especially in the southern region. Up to 65 feet of snow may accumulate on the high mountains of this province. At

the lowest elevations of the Cascade Province, vegetation is incredibly dense, and the conifer forests are comprised of Douglas fir, western redcedar, western hemlock, grand fir, Sitka spruce, and Alaska cedar. In the humid forests of southwestern Oregon, silver fir and redwood are dominant.

Long before there were homes and farms in this area, the original, natural vegetation provided continuous cover and adjacent feeding opportunities for wildlife, including pollinators. In choosing plants, aim to create habitat for pollinators that allow adequate food shelter, and water sources. Most pollinators have very small home ranges. You can make a difference by understanding the vegetation patterns of the farm, forest, or neighbor's yard adjacent to you and by making planting choices that support the pollinators' need for food and shelter as they move through the landscape.


# UNDERSTANDING THE CASCADE MIXED FOREST PROVINCE



- ✿ This region is designated **number M242** in the Baileys' Ecosystem Provinces. To see a map of the provinces go to: [www.fs.fed.us/colorimagemap/ecoreg1\\_provinces.html](http://www.fs.fed.us/colorimagemap/ecoreg1_provinces.html)
- ✿ Not sure about which bioregion you live or work in? Go to [www.pollinator.org](http://www.pollinator.org) and click on **Ecoregion Locator** for help.
- ✿ 53,400 square miles within Oregon and Washington.
- ✿ Primarily mountainous with some volcanoes.
- ✿ Elevations ranging from sea level to over 14,000 feet.
- ✿ Average annual temperature range from 35° to 50°F.
- ✿ Average year-round precipitation between 30-150 inches.
- ✿ USDA Hardiness Zones 8b-5a (1990 version).

## CHARACTERISTICS

- ✿ Dominated by Douglas fir, western redcedar, western hemlock, grand fir, Sitka spruce, and Alaska cedar at lowest elevations.
- ✿ Dominated by silver fir and redwood in humid forests of southwest Oregon.
- ✿ Ponderosa pine forests on dry eastern slopes of the Cascades.
- ✿ All but the highest peaks covered in vegetation.



The Cascade Mixed Forest, Coniferous Forest, Alpine Meadow Province includes:

- Oregon
- Washington

**“ ADDING NATIVE PLANTINGS IN RIPARIAN AREAS TO IMPROVE POLLINATOR HABITAT MAKES SENSE IN ADVANCING OUR FAMILY FARM’S CONSERVATION AND ECONOMIC OBJECTIVES, ENHANCING BENEFICIAL WILDLIFE AND IMPROVING POLLINATION IN OUR ORCHARD AND GARDEN. ”**

—LEE MCDANIEL, FARMER AND PRESIDENT, NATIONAL ASSOCIATION OF CONSERVATION DISTRICTS

# MEET THE POLLINATORS



Flower flies, also known as hoverflies, mimic the appearance of bees or wasps and feed on nectar and pollen like bees, but are classified as flies.

Western Tiger Swallowtail feeding on nectar in Olympia, Washington.



## WHO ARE THE POLLINATORS?

### BEES

Bees are well documented pollinators in the natural and agricultural systems of the Cascade Mixed Forest. A wide range of crops including apples, broccoli and cranberries are just a few plants that benefit from bee pollinators.

Most of us are familiar with the colonies of honey bees that have been the workhorses of agricultural pollination for years in the United States. They were imported from Europe almost 400 years ago.

There are nearly 4000 species of native ground and twig nesting bees in the U.S. Some form colonies while others live and work a solitary life. Native bees currently pollinate many crops and can be encouraged to do more to support agricultural endeavors if their needs for nesting habitat are met and if suitable sources of nectar, pollen, and water are provided. Bees have tongues of varying lengths that help determine which flowers they can obtain nectar and pollen from.

The bumble bee (*Bombus* spp.) forms small colonies, usually underground. They are generalists, feeding on a wide range of plant material from February to November and are important pollinators of tomatoes. The sweat bee (family *Halictidae*) nests underground. Various species are solitary while others form loose colonies.

Solitary bees include carpenter bees (*Xylocopa* spp.), which nest in wood; digger, or polyester bees (*Colletes* spp.), which nest underground; leafcutter bees (*Megachile* spp.), which prefer dead trees or branches for their nest sites; and mason bees (*Osmia* spp.), which utilize cavities that they find in stems and dead wood. Cactus bees (*Diadasia* spp.) are also solitary ground nesters.

### BUTTERFLIES

Gardeners have been attracting butterflies to their gardens for some time. These insects tend to be eye-catching, as are the flowers that attract them. Position flowering plants where they have full sun and are protected from the wind. Also, you will need to provide open areas (e.g. bare earth, large stones) where butterflies may bask, and moist soil from which they may get needed minerals. By providing a safe place to eat and nest, gardeners can also support the pollination role that butterflies play in the landscape. It might mean accepting slight damage to the plants, known as host plants, that provide food for the larval stage of the butterfly.

A diverse group of butterflies are present in garden areas and woodland edges that provide bright flowers, water sources, and specific host plants. Numerous trees, shrubs, and herbaceous plants support butterfly populations.

Butterflies are in the Order *Lepidoptera*. Some of the species in the Cascade Mixed Forest are



Brush-footed, Gossamer-winged, Swallowtail, Parnassian, Skipper, White, Sulphur and Milkweed butterflies. They usually look for flowers that provide a good landing platform.

Wet mud areas provide butterflies with both the moisture and minerals they need to stay healthy. Butterflies eat rotten fruit and even dung, so don't clean up all the messes in your garden!

## MOTHS

Moths are most easily distinguished from butterflies by their antennae. Butterfly antennae are simple with a swelling at the end. Moth antennae differ from simple to featherlike, but never have a swelling at the tip. In addition, butterflies typically are active during the day; moths at night. Butterfly bodies are not very hairy, while moth bodies are quite hairy and more stout.

Moths, generally less colorful than butterflies, also play a role in pollination. They are attracted to flowers that are strongly sweet smelling, open in late afternoon or night, and are typically white or pale colored.

## BEETLES

Over 30,000 species of beetles are found in the United States and many of them can be found on flower heads. Gardeners have yet to intentionally draw beetles to their gardens, possibly because beetle watching isn't as inspiring

as butterfly or bird watching. Yet beetles do play a role in pollination. Some have a bad reputation because they can leave a mess behind, damaging plant parts that they eat. Beetles are not as efficient as some pollinators. They wander between different species, often dropping pollen as they go.

Beetle pollinated plants tend to be large, strong scented flowers with their sexual organs exposed. They are known to pollinate Magnolia, sweetshrub (*Calycanthus*), paw paws, and yellow pond lilies.

## FLIES

It may be hard to imagine why one would want to attract flies to the garden. However, like beetles, the number of fly species and the fact that flies are generalist pollinators (visit many species of plants), should encourage us all to leave those flies alone and let them do their job as pollinators.

Recent research indicates that flies primarily pollinate small flowers that bloom under shade and in seasonally moist habitats. The National Research Council's *Status of Pollinators in North America* study states that flies are economically important as pollinators for a range of annual and bulbous ornamental flowers.

Plants pollinated by the fly include the American pawpaw (*Asimina triloba*), dead horse arum (*Helicodiceros muscivorus*), skunk cabbage (*Symplocarpus foetidus*), goldenrod (*Solidago* spp.), and

members of the carrot family like Queen Anne's lace (*Daucus carota*).

## BIRDS

Hummingbirds are the primary birds which play a role in pollination in North America. Their long beaks and tongues draw nectar from tubular flowers. Pollen is carried on both the beaks and feathers of different hummingbirds. The regions closer to the tropics, with warmer climates, boast the largest number of hummingbird species and the greatest number of native plants to support the bird's need for food. White-winged doves (*Zenaidura asiatica*) are also pollinators of the saguaro cactus (*Carnegieia gigantea*) in the south central United States.

Bright colored tubular flowers attract hummingbirds to gardens throughout the United States. Hummingbirds can see the color red; bees cannot. Many tropical flowers, grown as annuals in the Cascade Mixed Forest, along with native woodland edge plants, attract hummingbirds.

## BATS

Though bats in the Cascade Mixed Forest are not pollinators, bats play an important role in pollination in the southwest where they feed on agave and cactus. The long-nosed bats' head shape and long tongue allows it to delve into flower blossoms and extract both pollen and nectar.

## PLANT TRAITS

### WHICH FLOWERS DO THE POLLINATORS PREFER?

NOT ALL POLLINATORS ARE found in each North American province, and some are more important in different parts of the United States. Use this page as a resource to understand the plants and pollinators where you live.

Plants can be grouped together based on the similar characteristics of their flowers. These floral characteristics can be useful to predict the type of pollination method or animal that is most effective for that group of plants. This association between floral characteristics and pollination method is called a pollination syndrome.

The interactions of animal pollinators and plants have influenced the evolution of both groups of organisms. A mutualistic relationship between the pollinator and the plant species helps the pollinator find necessary pollen and nectar sources and helps the plant reproduce by ensuring that pollen is carried from one flower to another.

Plant Trait	Bats	Bees	Beetles
Color	Dull white, green or purple	Bright white, yellow, blue, or UV	Dull white or green
Nectar guides	Absent	Present	Absent
Odor	Strong musty; emitted at night	Fresh, mild, pleasant	None to strongly fruity or fetid
Nectar	Abundant; somewhat hidden	Usually present	Sometimes present; not hidden
Pollen	Ample	Limited; often sticky and scented	Ample
Flower Shape	Regular; bowl shaped – closed during day	Shallow; have landing platform; tubular	Large bowl-like, Magnolia

This chart and more information on pollinator syndromes can be found at:



## AND THE POLLINATORS THEY ATTRACT

### Pollinator

Birds	Butterflies	Flies	Moths	Wind
Scarlet, orange, red or white	Bright, including red and purple	Pale and dull to dark brown or purple; flecked with translucent patches	Pale and dull red, purple, pink or white	Dull green, brown, or colorless; petals absent or reduced
Absent	Present	Absent	Absent	Absent
None	Faint but fresh	Putrid	Strong sweet; emitted at night	None
Ample; deeply hidden	Ample; deeply hidden	Usually absent	Ample; deeply hidden	None
Modest	Limited	Modest in amount	Limited	Abundant; small, smooth, and not sticky
Large funnel like; cups, strong perch support	Narrow tube with spur; wide landing pad	Shallow; funnel like or complex and trap-like	Regular; tubular without a lip	Regular; small and stigmas exerted

<http://www.fs.fed.us/wildflowers/pollinators/syndromes.shtml>



# DEVELOPING LANDSCAPE PLANTINGS THAT PROVIDE POLLINATOR HABITAT

WHETHER YOU ARE A FARMER of many acres, land manager of a large tract of land, or a gardener with a small lot, you can increase the number of pollinators in your area by making conscious choices to include plants that provide essential habitat for bees, butterflies, moths, beetles, hummingbirds and other pollinators.

## FOOD:

Flowers provide nectar (high in sugar and necessary amino acids) and pollen (high in protein) to pollinators.

Fermenting fallen fruits also provide food for bees, beetles and butterflies.

Specific plants, known as host plants, are eaten by the larvae of pollinators such as butterflies.

- Plant in groups to increase pollination efficiency. If a pollinator can visit the same type of flower over and over, it doesn't have to relearn how to enter the flower and can transfer pollen to the same species, instead of squandering the pollen on unreceptive flowers.
- Plant with bloom season in mind, providing food from early spring to late fall. (see Bloom Periods pp.16-17)
- Plant a diversity of plants to support a variety of pollinators. Flowers of different color, fragrance, and season of bloom on plants of different heights will attract different pollinator species and provide pollen and nectar throughout the seasons.
- Many herbs and annuals, although

not native, are very good for pollinators. Mint, oregano, garlic, chives, parsley and lavender are just a few herbs that can be planted. Old fashioned zinnias, cosmos, and single sunflowers support bees and butterflies.

- Recognize weeds that might be a good source of food. For example, dandelions provide nectar in the early spring before other flowers open. Plantain is alternate host for the Baltimore Checkerspot.
- Learn and utilize Integrated Pest Management (IPM) practices to address pest concerns. Minimize or eliminate the use of pesticides.

## SHELTER:

Pollinators need protection from severe weather and from predators as well as sites for nesting and roosting.

- Incorporate different canopy layers in the landscape by planting trees, shrubs, and different-sized perennial plants.
- Leave dead snags for nesting sites of bees, and other dead plants and leaf litter for shelter.
- Build bee boxes to encourage solitary, non-aggressive bees to nest on your property.
- Leave some areas of soil uncovered to provide ground nesting insects easy access to underground tunnels.
- Group plantings so that pollinators can move safely through the landscape protected from predators.
- Include plants that are needed

by butterflies during their larval development.

## WATER:

A clean, reliable source of water is essential to pollinators.

- Natural and human-made water features such as running water, pools, ponds, and small containers of water provide drinking and bathing opportunities for pollinators.
- Ensure the water sources have a shallow or sloping side so the pollinators can easily approach the water without drowning.

Your current landscape probably includes many of these elements. Observe wildlife activity in your farm fields, woodlands, and gardens to determine what actions you can take to encourage other pollinators to feed and nest. Evaluate the placement of individual plants and water sources and use your knowledge of specific pollinator needs to guide your choice and placement of additional plants and other habitat elements. Minor changes by many individuals can positively impact the pollinator populations in your area. Watch for - and enjoy - the changes in your landscape!

- **CAUTION:** Remember that pesticides are largely toxic to pollinators. Extreme caution is warranted if you choose to use any pesticide. Strategically apply pesticides only for problematic target species.



# FARMS

Raspberries, alfalfa, cucumbers, apples and strawberries are a few of the food crops in the Cascade Mixed Forest Province that will benefit from strong native bee populations that boost pollination efficiency. Incorporate different plants throughout the farm that provide food for native populations when targeted crops are not in flower.

Farmers have many opportunities to incorporate pollinator-friendly land management practices on their land which will benefit the farmer in achieving his or her production goals:

- Manage the use of pesticides to reduce the impact on native pollinators. Spray when bees aren't active (just after dawn) and choose targeted ingredients.
- Carefully consider the use of

herbicides. Perhaps the targeted weeds can provide needed food for pollinators.

- Minimize tillage to protect ground nesting pollinators.
- Ensure water sources are scattered throughout the landscape.
- Choose a variety of native plants to act as windbreaks, riparian buffers, and field borders throughout the farm.
- Plant unused areas of the farm with temporary cover crops that can provide food or with a variety of trees, shrubs, and flowers that provide both food and shelter for pollinators.
- Check with your local Natural Resources Conservation Service (NRCS) office to see what technical and financial support might be available to assist you in your effort to provide nectar, pollen, and larval food sources for pollinators on your farm.

“**FOOD SUPPLIES FOR BEES ARE CRITICAL TO MAINTAINING STRONG HIVES FOR ALMOND POLLINATION THE FOLLOWING WINTER.**”

-- DAN CUMMINGS,  
CHICO, CALIFORNIA  
ALMOND GROWER.



Illustrations by Carolyn Vibbert

## PUBLIC LANDS

“FROM HUMMINGBIRDS TO BEETLES, TO BUTTERFLIES, NATURE’S POLLINATORS HELP KEEP MIDEWIN’S TALLGRASS PRAIRIE RESTORATIONS FULL OF DIVERSE FLOWERING PLANTS. INSECT MONITORING PROVIDES A KEY MEASURE OF OUR SUCCESS.”

-- LOGAN LEE  
PRAIRIE SUPERVISOR, MIDEWIN  
NATIONAL TALLGRASS PRAIRIE



Public lands are maintained for specific reasons ranging from high impact recreation to conservation. In the Cascade Mixed Forest, forests have been cut to allow for roads, buildings, open lawn areas, boat ramps, and vistas. Less disturbed natural areas can be augmented with plantings of native plant species. Existing plantings around buildings and parking areas should be evaluated to determine if pollinator-friendly plants can be substituted or added to attract and support pollinators. Public land managers have a unique opportunity to use their plantings as an education tool to help others understand the importance of pollinators in the environment through signs, brochures, and public programs.

In an effort to increase populations of pollinators the land manager can:

- Inventory and become knowledgeable of local pollinators.
- Provide connectivity between vegetation areas by creating corridors of perennials, shrubs, and trees that provide pollinators shelter and food as they move through the landscape.
- Maintain a minimum of lawn areas that support recreational needs.
- Restrict the use of pesticides and herbicides.
- Provide water sources in large open areas.
- Maintain natural meadows and openings that provide habitats for sun-loving wildflowers and grasses.
- Remove invasive species and encroaching shrubs and trees.



## HOME LANDSCAPES

**“ A GARDEN IS ONLY AS RICH AND BEAUTIFUL AS THE INTEGRAL HEALTH OF THE SYSTEM; POLLINATORS ARE ESSENTIAL TO THE SYSTEM - MAKE YOUR HOME THEIR HOME. ”**

-- DERRY MACBRIDE  
NATIONAL AFFAIRS AND  
LEGISLATION CHAIRWOMAN,  
GARDEN CLUB OF AMERICA

Gardeners have a wide array of plants to use in their gardens. Native plants, plants introduced from years of plant exploration from around the world, and plants developed by professional and amateur breeders can be found in garden centers, in catalogs, and on web-sites. Use your knowledge of pollinator needs to guide your choices.

- Choose a variety of plants that will provide nectar and pollen throughout the growing season.
- Resist the urge to have a totally manicured lawn and garden. Leave bare ground for ground nesting bees. Leave areas of dead wood and leaf litter for other insects.
- Strive to eliminate the use of all pesticides.
- Find local resources to help you in your efforts. Contact your local county extension agent or native plant society. Visit your regional botanic gardens and arboreta.

The scale of your plantings will vary but it is important to remember that you are trying to provide connectivity to the landscape adjacent to your property. Don't just look within your property boundaries. If your neighbor's property provides an essential element, such as water, which can be utilized by pollinators visiting your land, you may be able to devote more space to habitat elements that are missing nearby. It is best to use native plants which have evolved to support the needs of specific native pollinators. Some pollinators, however, are generalists and visit many different plants, both native and non-native. Be sure that any non-native plants you choose to use are not invasive. Remember that specialized cultivars sometimes aren't used by pollinators. Flowers that have been drastically altered, such as those that are double or a completely different color than the wild species, often prevent pollinators from finding and feeding on the flowers. In addition, some altered plants don't contain the same nectar and pollen resources that attract pollinators to the wild types.

- **CAUTION:** Take time to evaluate the source of your plant material. You want to ensure you get plants that are healthy and correctly identified. Your local native plant society can help you make informed decisions when searching for plants.



# BLOOM PERIODS

## FOR THE CASCADE MIXED FOREST PROVINCE

The following chart lists plants and the time they are in bloom throughout the growing seasons. Choose a variety of flower colors and make sure something is blooming at all times! Note for all charts: When more than one species of the same genus is useful, the genus name is followed by “spp.”

Botanical Name	Common Name	Habitat	Feb	March	April	May	June	July	Aug	Sept	Oct
<b>Trees &amp; Shrubs</b>											
<i>Acer circinatum</i>	vine maple	Forest		red	red	red	red				
<i>Acer macrophyllum</i>	big-leaf maple	Forest		greenish white	greenish white	greenish white	greenish white				
<i>Amelanchier alnifolia</i>	serviceberry	Forest			white	white	white	white			
<i>Arbutus menziesii</i>	madrone	Forest			white	white					
<i>Arctostaphylos uva-ursi</i>	kinnikinnick	Forest			white	white	white				
<i>Berberis aquifolium</i>	tall Oregon grape	Forest		yellow	yellow	yellow					
<i>Berberis nervosa</i>	dwarf Oregon grape	Forest		yellow	yellow	yellow	yellow				
<i>Calocedrus decurrens</i>	incense cedar	Forest		cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing
<i>Cassiope mertensiana</i>	Merten's mountain heather	subalpine/ alpine meadow						white	white		
<i>Cornus nuttallii</i>	Pacific dogwood	Forest			white	white	white				
<i>Gaultheria shallon</i>	salal	Forest				white to pink	white to pink	white to pink			
<i>Holodiscus discolor</i>	ocean-spray	Forest					white	white	white		
<i>Luetkea pectinata</i>	Luetkea	subalpine/ alpine meadow					white	white	white		
<i>Phyllodoce empetriformis</i>	pink mountain heather	subalpine/ alpine meadow					pinkish-rose	pinkish-rose	pinkish-rose		
<i>Picea sitchensis</i>	Sitka spruce	Forest		cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing
<i>Pinus lambertiana</i>	sugar pine	Forest		cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing
<i>Pseudotsuga menziesii</i>	Douglas-fir	Forest		cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing
<i>Rhododendron macrophyllum</i>	rhododendron	Forest				pink	pink	pink			
<i>Rosa gymnocarpa</i>	baldhip rose	Forest					pink	pink			
<i>Rubus parviflorus</i>	thimbleberry	Forest				white	white	white			
<i>Rubus spectabilis</i>	salmonberry	Forest		deep pink	deep pink	deep pink	deep pink				
<i>Saxifraga tolmiei</i>	Tolmie's saxifrage	subalpine/ alpine meadow						white	white		
<i>Symporicarpos alba</i>	snowberry	Forest					pink	pink			
<i>Thuja plicata</i>	western red cedar	Forest		cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing
<i>Tsuga heterophylla</i>	western hemlock	Forest		cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing	cone bearing
<i>Vaccinium deliciosum</i>	blue huckleberry	subalpine/ alpine meadow				pink	pink				
<i>Vaccinium membranaceum</i>	big huckleberry	Forest			yellowish pink	yellowish pink	yellowish pink				
<i>Vaccinium ovatum</i>	evergreen huckleberry	Forest			pink	pink	pink	pink	pink		
<i>Vaccinium parvifolium</i>	red huckleberry	Forest			pale pink	pale pink	pale pink				
<b>Perennial Flowers</b>											
<i>Achlys triphylla</i>	vanilla leaf	Forest			white	white	white	white			
<i>Anemone occidentalis</i>	mountain pasque flower	subalpine/ alpine meadow				white to purple tinged					
<i>Aquilegia formosa</i>	Cascade columbine	Forest				red	red	red	red		
<i>Arnica latifolia</i>	broadleaf arnica	Forest					yellow	yellow			

Botanical Name	Common Name	Habitat	Feb	March	April	May	June	July	Aug	Sept	Oct
<b>Perennial Flowers continued</b>											
<i>Asarum caudatum</i>	wild ginger	Forest			reddish green	reddish green	reddish green				
<i>Castilleja parviflora</i>	mountain indian paintbrush	subalpine/ alpine meadow					rose to crimson	rose to crimson	rose to crimson		
<i>Clintonia uniflora</i>	queencup beadlily	Forest					white	white			
<i>Cornus canadensis</i>	bunchberry dogwood	Forest					white	white	white		
<i>Dicentra formosa</i>	Pacific bleeding heart	Forest		pink	pink	pink	pink	pink			
<i>Chamerion angustifolium</i> spp. <i>angustifolium</i>	fireweed	subalpine/ alpine meadow					rose to purple	rose to purple	rose to purple	rose to purple	
<i>Erigeron peregrinus</i>	subalpine fleabane	subalpine/ alpine meadow						white to rose purple	white to rose purple		
<i>Erythronium oregonum</i>	giant fawnlily	Forest		white/yellow base	white/yellow base	white/yellow base					
<i>Iris tenax</i>	Oregon iris	Forest			purple	purple	purple				
<i>Lilium columbianum</i>	Columbia lily	Forest					yellow orange to red orange	yellow orange to red orange			
<i>Lilium washingtonianum</i>	Washington lily	Forest					white to pale pink	white to pale pink			
<i>Linnaea borealis</i>	twinflower	Forest					pink	pink	pink	pink	
<i>Lupinus latifolius</i>	broadleaf lupine	Forest					blue to purple	blue to purple	blue to purple		
<i>Lysichitum americanum</i>	skunk cabbage	Forest	yellow	yellow	yellow	yellow	yellow	yellow			
<i>Maianthemum dilatatum</i>	false lily of the valley	Forest				white	white				
<i>Mimulus lewisii</i>	monkey flower	subalpine/ alpine meadow					pink-purple	pink-purple	pink-purple		
<i>Oreostemma alpigenum</i>	tundra aster	subalpine/ alpine meadow						violet to lavender	violet to lavender		
<i>Oxalis oregana</i>	Oregon oxalis	Forest			white to pink	white to pink	white to pink	white to pink	white to pink	white to pink	
<i>Pedicularis groenlandica</i>	elephant's head	subalpine/ alpine meadow					pink-purple	pink-purple	pink-purple		
<i>Polygonum bistortoides</i>	bistort	subalpine/ alpine meadow				white to pinkish	white to pinkish	white to pinkish	white to pinkish		
<i>Potentilla gracilis</i> var. <i>flabelliformis</i>	high mountain cinquefoil	subalpine/ alpine meadow					yellow	yellow	yellow		
<i>Prosartes hookeri</i> var. <i>hookeri</i>	hooker fairybells	Forest			greenish white	greenish white	greenish white	greenish white			
<i>Prosartes smithii</i>	Smith fairybells	Forest			greenish white	greenish white	greenish white	greenish white			
<i>Satureja douglasii</i>	yerba buena	Forest					white to pink	white to pink			
<i>Maianthemum racemosum</i> ssp. <i>racemosum</i>	false solomonplume	Forest			white	white	white	white			
<i>Streptopus amplexifolius</i>	twisted-stalk	Forest				white	white	white			
<i>Trillium albidum</i>	trillium	Forest		white to purple	white to purple	white to purple	white to purple				
<i>Trillium ovatum</i>	trillium	Forest		white to purple	white to purple	white to purple	white to purple				
<i>Valeriana sitchensis</i>	Sitka valerian	Forest					white to pink	white to pink	white to pink		
<i>Vancouveria hexandra</i>	inside-out-flower	Forest				white	white				
<i>Veratrum viride</i>	false hellebore	subalpine/ alpine meadow					yellow-green to green	yellow-green to green	yellow-green to green	yellow-green to green	
<i>Viola sempervirens</i>	redwoods violet	Forest		yellow	yellow	yellow	yellow				
<i>Xerophyllum tenax</i>	beargrass	Forest				white	white	white	white		
<b>Vines</b>											
<i>Lonicera hispidula</i>	hairy honeysuckle	Forest					pink, yellowish pink	pink, yellowish pink	pink, yellowish pink		
<i>Rubus ursinus</i>	trailing blackberry	Forest			white	white	white	white	white		
<i>Whipplea modesta</i>	whipple vine	Forest			white	white	white				

# PLANTS THAT ATTRACT POLLINATORS

## IN THE CASCADE MIXED FOREST PROVINCE

The following chart lists plants that attract pollinators. It is not exhaustive, but provides guidance on where to start. Annuals, herbs, weeds, and cover crops provide food and shelter for pollinators, too.

Botanical Name	Common Name	Color	Height (m)	Flower Season	Sun	Soil	Visitation by Pollinator	Also a host plant
<b>Trees &amp; Shrubs</b>								
<i>Acer</i> spp.	maple	greenish white to red	< 30	March–June	sun to partial shade	moist, well drained	bees	X
<i>Amelanchier alnifolia</i>	serviceberry	white	1 - 5	April–July	sun to partial shade	moist to dry	bees, flies	X
<i>Arbutus menziesii</i>	madrone	white	6 - 30	April–May	sun to partial shade	dry	bees	
<i>Arctostaphylos</i> spp.	kinnikinnick	white	0.1 - 4	April–July	sun to partial shade	dry, well drained	hummingbirds	
<i>Cornus nuttallii</i>	Pacific dogwood	white	1 - 30	April–June	shade	moist, well drained	bees, beetles, flies, butterflies	
<i>Rhododendron macrophyllum</i>	rhododendron	pink	1 - 5	May–July	shade	moist, well drained	bees	
<i>Ribes</i> spp.	currants/ gooseberrys	greenish white, white, pink, red	1 - 3	March–June	sun to shade	moist to dry, well drained	hummingbirds	
<i>Sambucus</i> spp.	elderberry	white to creamy	1 - 6	May–July	sun to partial shade	moist to dry, well drained	bees	
<i>Symphoricarpos</i> spp.	snowberry	pink	0.5 - 2	May–Aug	sun to shade	moist, well drained	bees	X
<i>Vaccinium</i> spp.	huckleberry	pink	0.1 - 3	April–Aug	sun to partial shade	moist to dry, well drained	bees	
<b>Perennial Flowers</b>								
<i>Achillea millefolium</i>	yarrow	white	0.2 - 1	April–Oct	sun to partial shade	dry	bees	X
<i>Aquilegia formosa</i>	Cascade columbine	red	0.1 - 1	May–Aug	partial shade to shade	moist	hummingbirds, bees	
<i>Aster</i> spp.	daisy	blue, purple, pink to white	0.1 - 1	July–Sept	sun to partial shade	moist to dry	bees	
<i>Delphinium</i> spp.	larkspur	white to blue	0.1 - 0.5	April–July	sun to partial shade	dry	bees	



Botanical Name	Common Name	Color	Height (m)	Flower Season	Sun	Soil	Visitation by Pollinator	Also a host plant
<b>Perennial Flowers continued</b>								
<i>Erigeron</i> spp.	fleabane	blue, purple, pink to white	0.1 - 0.7	June–Aug	sun to partial shade	moist to dry	bees, butterflies, moths	
<i>Eriogonum</i> spp.	buckwheat	white to yellow	0.1 - 0.5	May–Aug	sun	dry, well drained	bees	
<i>Erythronium</i> spp.	fawnlily	white, pink, yellow	0.1 - 0.3	March–Aug	sun to shade		bees	
<i>Eschscholzia californica</i>	California poppy	yellow to orange	0.1 - 0.5	May–Sept	sun	dry, well drained	bees	
<i>Hydrophyllum</i> spp.	waterleaf	white, blue, purple	0.2 - 0.8	April–July	sun to shade	moist	bees	
<i>Iris</i> spp.	iris	white, yellow to purple	0.1 - 0.4	April–July	sun to partial shade		bees	
<i>Lilium</i> spp.	lily	white to pinkish, orange	0.2 - 1	June–July	sun to partial shade	moist	hummingbirds	
<i>Lupinus</i> spp.	lupine	blue to purple	0.1 - 1	April–Aug	sun to partial shade	dry to moist	bees	X
<i>Mentha arvensis</i>	mint	white to pink or purpleq	0.2 - 0.8	July–Sept	sun to partial shade	moist	bees	
<i>Penstemon</i> spp.	penstemon	white to purple or red	0.1 - 0.6	May–Aug	sun to partial shade	dry, well drained	bees	X
<i>Phacelia</i> spp.	scorpion weed	white	0.2 - 1	May–Aug	sun	dry, well drained	bees	
<i>Sedum</i> spp.	stonecrop	white, pink, yellow	0.1 - 0.3	May–Aug	sun	dry, well drained	bees	X
<i>Solidago</i> spp.	goldenrod	yellow	0.3 - 2	July–Oct	sun to partial shade	moist	bees, butterflies, beetles, wasps	X
<i>Trillium</i> spp.	trillium	white to purple	0.1 - 0.3	March–June	partial shade to shade		beetles, flies, bees	
<b>Vines</b>								
<i>Lonicera hispidula</i>	hairy honeysuckle	pink, yellowish pink	< 6	June - Aug	partial shade to shade	dry to moist	hummingbirds	

**References:** Franklin, J. F. and Dyrness, C. T. 1973. Natural vegetation of Oregon and Washington. Gen. Tech. Rep. PNW 8. USDA Forest Service, Portland, OR, 417p. • Hitchcock, C. L. and Cronquist, A. 1973. Flora of the Pacific Northwest. University of Washington Press, Seattle, WA, 730p. • North American Pollinator Protection Campaign and the Coevolution Institute. 2007. Hummers. <http://www.pollinator.org/Resources/Hummers.pdf> • Peck, M. 1961. A Manual of higher plants of Oregon. Binforids & Mort Publishers, Portland, OR, 936 p. • Rose, R.; Chachulski, C. E. C.; Haase, D. L. 1998. Propagation of Pacific Northwest native plants. Oregon State University Press, Corvallis, OR, 246p. • USDA Natural Resources Conservation Service. 2007. Plants database. <http://plants.usda.gov/index.html> Xerces Society for Invertebrate Conservation. 2007. Butterfly gardening. [http://www.xerces.org/Pollinator\\_Insect\\_Conservation/Xerces\\_butterfly\\_gardening\\_fact\\_sheet.pdf](http://www.xerces.org/Pollinator_Insect_Conservation/Xerces_butterfly_gardening_fact_sheet.pdf) Xerces Society for Invertebrate Conservation. 2007. Plants for bees in the Pacific Northwest. [http://www.xerces.org/Pollinator\\_Insect\\_Conservation/Xerces\\_PNW\\_plants\\_fact\\_sheet.pdf](http://www.xerces.org/Pollinator_Insect_Conservation/Xerces_PNW_plants_fact_sheet.pdf)

# HABITAT HINTS

## FOR THE CASCADE MIXED FOREST PROVINCE

### HABITAT REQUIREMENTS FOR BEE-POLLINATED GARDEN FLOWERS AND CROPS

	Bumble	Digger	Lg Carpenter	Sm Carpenter	Squash/ Gourd	Leafcutter	Mason	Sweat	Plasterer	Yellow-faced	Andrenid
<b>FLOWERS</b>											
Catalpa			x								
Catnip	x	x					x				
Clover		x									x
Columbine	x										
Cow parsley										x	
Goldenrod	x	x				x		x			
Impatiens	x										
Irises	x		x								
Lavender	x	x	x			x					
Milkwort								x			
Morning glory				x							
Penstemon	x	x					x				
Passion flowers			x								
Phacelia	x	x		x		x	x	x	x		x
Potentilla										x	
Rose	x		x				x	x		x	
Salvia	x	x	x			x	x				
Saxifrages								x		x	
Sorrel				x							
Sunflowers	x	x	x	x		x		x	x		x
Violet								x			x
Wild Mustard		x							x		
Willow catkins									x		x
<b>CROPS</b>											
Almond	x						x				x
Apple							x				
Blueberry	x	x									x
Cherry							x				x
Eggplant	x		x					x			
Gooseberry	x										x
Legumes	x	x				x		x			
Water melon	x							x			
Squash/ Pumpkins/ Gourds			x		x						
Tomatoes	x	x	x					x			
Thyme	x	x					x	x		x	



## HABITAT AND NESTING REQUIREMENTS:

### **Bumble Bees:**

Abandoned mouse nests, other rodent burrows, upside down flower pots, under boards, and other human-made cavities. Colonies are founded by a queen in the spring and don't die out in the fall. New queens mate then and overwinter in a sort of hibernation. Bumble bees are usually active during the morning hours and forage at colder temperatures than honey bees, even flying in light rain.

### **Large carpenter bees:**

Soft dead wood, poplar, cottonwood or willow trunks and limbs, structural timbers including redwood. Depending on the species, there may be one or two brood cycles per year. These bees can be active all day even in the hottest weather.

### **Digger bees:**

Sandy soil, compacted soils, bank sides. Anthophorid bees (now in the Apidae) are usually active in the morning hours, but can be seen at other times.

### **Small carpenter bees:**

Pithy stems including roses and blackberry canes. These bees are more active in the morning but can be found at other times.

### **Squash and Gourd bees:**

Sandy soil, may nest in gardens (where pumpkins, squash and gourds are grown) or pathways. These bees are early risers and can be found in pumpkin patches before dawn. Males often sleep in the wilted flowers.

### **Leafcutter bees:**

Pre-existing circular tunnels of various diameters in dead but sound wood created by emerging beetles, some nest in the ground. Leave dead limbs and trees to support not just pollinators but other wildlife. Leafcutter bees can be seen foraging throughout the day even in hot weather.

### **Mason bees:**

Pre-existing tunnels, various diameters in dead wood made by emerging beetles, or human-made nesting substrates, drilled wood boards, paper soda straws inserted into cans attached to buildings. Mason bees are generally more active in the morning hours.

### **Sweat bees:**

Bare ground, compacted soil, sunny areas not covered by vegetation. Like most bees, sweat bees forage for pollen earlier in the morning and then for nectar later.

### **Plasterer or cellophane bees:**

Bare ground, banks or cliffs. Colletid bees can be active in the morning or later in the day.

### **Yellow-faced bees:**

In dead stems. These bees are more active during morning hours.

### **Andrenid bees:**

Sunny, bare ground, sand soil, under leaf litter or in soil in banksides and cliffs. These generally spring-active bees are most commonly seen on flowers during the morning when pollen and nectar resources are abundant.



**“MONARCH  
BUTTERFLIES  
NEVER FAIL TO  
CATCH THE  
VISITOR’S EYE  
AND ALWAYS  
LEAD TO  
A TEACHABLE  
MOMENT.”**

-- LOGAN LEE,  
PRAIRIE SUPERVISOR  
MIDWIN NATIONAL  
TALLGRASS PRAIRIE

### **BECOME FAMILIAR WITH POLLINATORS IN YOUR LANDSCAPE.**

- ✿ Watch for activity throughout the day and the seasons.
- ✿ Keep a simple notebook of when and what comes to your garden.  
NOTE: It is not necessary to identify each species when you first get started. Simply note if it is a bee that likes the yellow flower that blooms in the fall.
- ✿ Consult a local field guide or web site when you are ready to learn more details.

### **ADD NATIVE PLANTS TO ATTRACT MORE NATIVE POLLINATORS.**

- ✿ List the plants you currently have in your landscape.
- ✿ Determine when you need additional flowers to provide nectar and pollen throughout the growing season.
- ✿ Add plants that provide additional seasons of bloom, create variable heights for shelter, and attract the types of pollinators you want.
- ✿ Don't forget to include host plants that provide food and shelter for larval development.
- ✿ Contact your local native plant society or extension agent for more help.

### **USE POLLINATOR FRIENDLY LANDSCAPE PRACTICES TO SUPPORT THE POLLINATORS YOU ATTRACT.**

- ✿ Use Integrated Pest Management Practices to address pest concerns.
- ✿ Tolerate a little mess – leave dead snags and leaf litter, keep areas bare for ground nesting insects, and leave some weeds that provide food for pollinators.
- ✿ Provide safe access to clean water.

### **NOTICE THE CHANGES THAT YOU HAVE HELPED TO CREATE!**



## RESOURCES

Many books, websites, and people were consulted to gather information for this guide. Use this list as a starting point to learn more about pollinators and plants in your area.

### BAILEY'S ECOREGION MAPS

#### USDA Forest Service

[http://www.fs.fed.us/land/ecosysmgmt/ecoreg1\\_home.html](http://www.fs.fed.us/land/ecosysmgmt/ecoreg1_home.html)

### POLLINATION/POLLINATORS

#### Pollinator Partnership

[www.pollinator.org](http://www.pollinator.org)

#### Coevolution Institute

[www.coevolution.org](http://www.coevolution.org)

#### Natural Resources Conservation Service

[www.nrcs.usda.gov](http://www.nrcs.usda.gov)

#### North American Pollinator Protection Campaign

[www.nappc.org](http://www.nappc.org)

#### USDA Forest Service

[www.fs.fed.us/wildflowers/pollinators/](http://www.fs.fed.us/wildflowers/pollinators/)

#### Wild Farm Alliance

[www.wildfarmalliance.org](http://www.wildfarmalliance.org)

#### The Xerces Society

[www.xerces.org](http://www.xerces.org)

#### Illinois Natural History Survey

[www.inhs.uiuc.edu](http://www.inhs.uiuc.edu)

Buchmann, S.L. and G.P. Nabhan. 1997. *The Forgotten Pollinators*. Island Press: Washington, DC.

Committee on the Status of Pollinators in North America. 2007. *Status of Pollinators in North America*. The National Academies Press: Washington, DC.

### NATIVE PLANTS

#### Plant Conservation Alliance

[www.nps.gov/plants](http://www.nps.gov/plants)

#### Seeds of Success

[www.nps.gov/plants/sos](http://www.nps.gov/plants/sos)

#### Lady Bird Johnson Wildflower Center

[www.wildflower.org/plants/](http://www.wildflower.org/plants/)

#### USDA Hardiness Zone Map

[www.usna.usda/Hardzone/](http://www.usna.usda/Hardzone/)

#### U.S. National Arboretum

[www.usna.usda.gov/Hardzone/ushzmap.html](http://www.usna.usda.gov/Hardzone/ushzmap.html)

USDA, NRCS. 2007. The PLANTS Database

[www.plants.usda.gov](http://www.plants.usda.gov), 19 July, 2007  
National Plant Data Center,  
Baton Rouge, LA 70874-4490 USA

### NATIVE BEES

#### National Sustainable Information Service

"Alternative Pollinators: Native Bees" by Lane Greer, NCAT Agriculture Specialist, Published 1999, ATTRA Publication #IP126  
[www.attra.ncat.org/attra-pub/nativebee.html](http://www.attra.ncat.org/attra-pub/nativebee.html)

#### Agriculture Research Service

*Plants Attractive to Native Bees* table  
[www.ars.usda.gov/Research/docs.htm?docid=12052](http://www.ars.usda.gov/Research/docs.htm?docid=12052)

### BUTTERFLIES AND MOTHS

Opler, Paul A., Harry Pavulaan, Ray E. Stanford, Michael Pogue, coordinators. 2006. *Butterflies and Moths of North America*. Bozeman, MT: NBII Mountain Prairie Information Node.

[www.butterfliesandmoths.org/](http://www.butterfliesandmoths.org/)  
(Version 07192007)

Pyle, Robert Michael. 1981. *National Audubon Society Field Guide to Butterflies*. Alfred A. Knopf: New York, NY.

#### North American Butterfly Association

[www.naba.org](http://www.naba.org)

## FEEDBACK

We need your help to create better guides for other parts of North America. Please e-mail your input to [feedback@pollinator.org](mailto:feedback@pollinator.org) or fax to 415-362-3070.

- ✿ How will you use this guide?
- ✿ Do you find the directions clear? If not, please tell us what is unclear.
- ✿ Is there any information you feel is missing from the guide?
- ✿ Any other comments?

**THANK YOU  
FOR TAKING  
THE TIME TO HELP!**

**POLLINATOR  
PARTNERSHIP**



**NAPPC**



Plant Conservation Alliance



---

RESEARCH AND WRITING:

**ELIZABETH L. LEY  
STEPHEN BUCHMANN, PH.D.  
RUSSELL HOLMES  
KATHERINE MCGUIRE**

---

EDITORIAL:

**LAURIE DAVIES ADAMS  
AND LARRY STRITCH, PH.D.**

---

PRODUCTION SUPERVISION:

**KATHERINE MCGUIRE**

---

DESIGN:

**MARGUERITE MEYER**

---

CONCEPT REVIEW:

American Farm Bureau Federation, Ron Gaskell  
Bureau of Land Management, Peggy Olwell, Carol Spurrier,  
Mary Byrne, Mary Tisdale, Elizabeth Wooster  
National Garden Association, Susanne DeJohn  
Plant Conservation Alliance – Edward Fletcher, Jean Giblette,  
Mary Ann Lawler, Ron Smith  
Smithsonian Institute, Department of Botany,  
Gary Krupnick, Ph.D.  
USDA - CSREES, Greg Crosby, Ph.D., Leslie Gilbert, Ph.D.  
USDA - Forest Service, David Pivorunas, Larry Stritch, Ph.D.  
USDA - Natural Resource Conservation Service, Doug Holy,  
Hilda Diaz-Soltero  
USDOI - US Fish and Wildlife Service, Karen Anderson,  
Don MacLean, Patricia DeAngelis, Ph.D.  
USGS - Steve Hilburger, Elizabeth Sellers

---

PHOTO CONTRIBUTORS:

Diane Petit

---

ILLUSTRATIONS:

Carolyn Vibbert

---

For a copy of this brochure, or for another region, visit [www.pollinator.org](http://www.pollinator.org)

**The Pollinator Partnership™/North American Pollinator Protection Campaign**

423 Washington St., 5th Floor, San Francisco, CA 94111 – 415-362-1137

[www.pollinator.org](http://www.pollinator.org)  [www.nappc.org](http://www.nappc.org)

