

SELECTING

PLANTS

FOR

POLLINATORS



A REGIONAL GUIDE FOR FARMERS, LAND MANAGERS, AND GARDENERS IN THE

HAWAIIAN ISLANDS PROVINCE

INCLUDING THE TROPICAL MOIST FOREST, TROPICAL DRY FOREST, AND TROPICAL GRASSLAND

& SHRUBLAND

POLLINATOR PARTNERSHIP and NAPPC

TABLE OF CONTENTS

| WHY SUPPORT POLLINATORS? | 4 |
|---------------------------------|----|
| GETTING STARTED | 5 |
| HAWAIIAN ISLANDS PROVINCE | 6 |
| MEET THE POLLINATORS | 8 |
| PLANT TRAITS | 10 |
| DEVELOPING PLANTINGS | 12 |
| FARMS | 13 |
| PUBLIC LANDS | 14 |
| HOME LANDSCAPES | 15 |
| BLOOM PERIODS | 16 |
| PLANTS THAT ATTRACT POLLINATORS | 18 |
| HABITAT HINTS | 20 |
| CHECKLIST | 22 |
| RESOURCES AND FEEDBACK | 23 |

different regions in the United States. We welcome your feedback to assist us in making future guides useful. Please contact us at feedback@pollinator.org

This is one of several guides for

Cover photos: Butterfly & Eleilei Bay courtesy Karl Magnacca, Kauai mountains and taro fields courtesy Lauren Rusert.

SELECTING PLANTS FOR POLLINATORS

A REGIONAL GUIDE FOR FARMERS, LAND MANAGERS, AND GARDENERS

IN THE

ECOLOGICAL REGION OF THE

HAWAIIAN ISLANDS PROVINCE

INCLUDING THE

TROPICAL MOIST FOREST,

TROPICAL DRY FOREST,

AND

TROPICAL GRASSLAND & SHRUBLAND

A NAPPC AND POLLINATOR PARTNERSHIP PUBLICATION

This guide was funded by Robert Batinovich, the Greene Family, Ruth and Chuck Adams, and the Hawaii Department of Agriculture with oversight by the Pollinator Partnership (www.pollinator.org), in support of the North American Pollinator Protection Campaign (NAPPC–www.nappc.org).

WHY SUPPORT POLLINATORS?

IN THEIR 1996 BOOK, THE FORGOTTEN POLLINATORS, Buchmann and Nabhan estimated that animal pollinators are needed for the reproduction of 90% of flowering plants and one third of human food crops. Each of us depends on these industrious pollinators in a practical way to provide us with the wide range of foods we eat. In addition, pollinators are part of the intricate web that supports the biological diversity in natural ecosystems that helps sustain our quality of life.

Abundant and healthy populations of pollinators can improve fruit set and quality, and increase fruit size. In farming situations this increases production per acre. In the wild, biodiversity increases and wildlife food sources increase.

Macadamia nuts, avocados, watermelon, guava, and coffee are some of the crops that rely on honey bees and native bees for pollination. Domestic honey bees pollinate approximately \$10 billion worth of crops in the U.S. each year.

Unfortunately, the numbers of both native pollinators and domesticated bee populations are declining. They are threatened by habitat loss, disease, and the excessive and inappropriate use of pesticides. The loss of commercial bees to Colony Collapse Disorder (CCD) has highlighted how severe the issues of proper hive management are to reduce stresses caused by disease, pesticide use, insufficient nutrition, and transportation practices. Currently, the pollination services that the commercial beekeeping industry provides are receiving much needed research and conservation resources. The efforts to understand the threats to commercial bees should help us understand other pollinators and their roles in the environment as well.

It is imperative that we take immediate steps to help pollinator populations thrive. The beauty of the situation is that by supporting pollinators' need for habitat, we support our own needs for food and support diversity in the natural world.

Thank you for taking time to consult this guide. By adding plants to your landscape that provide food and shelter for pollinators throughout their active seasons and by adopting pollinator friendly landscape practices, you can make a difference to both the pollinators and the people that rely on them.

R

Val Dolcini President & CEO Pollinator Partnership

66 **FLOWERING PLANTS ACROSS WILD. FARMED** AND EVEN URBAN LANDSCAPES ACTUALLY FEED THE TERRESTRIAL WORLD. AND POLLINATORS ARE THE **GREAT CONNECTORS** WHO ENABLE THIS GIANT FOOD SYSTEM TO WORK FOR ALL WHO EAT... 99 **INCLUDING US.** -- ROGER LANG. CHAIRMAN.

POLLINATOR PARTNERSHIP

GETTING STARTED



THIS REGIONAL GUIDE IS just one in a series of plant selection tools designed to provide information on how individuals can influence pollinator populations through choices they make when they farm a plot of ground, manage large tracts of public land, or plant a garden. Each of us can have a positive impact by providing the essential habitat requirements for pollinators including food, water, shelter, and enough space to allow pollinators to raise their young.

Pollinators travel through the landscape without regard to property ownership or state boundaries. We've chosen to use R.G. Bailey's classification system to identify the geographic focus of this guide and to underscore the connections between climate and vegetation types that affect the diversity of pollinators in the environment.

Bailey's Ecoregions of the United States, developed by the United States Forest Service, is a system created as a management tool and is used to predict responses to land management practices throughout large areas. This guide addresses pollinator-friendly land management practices in what is known as the Hawaiian Islands Province.

The Hawaiian Island Province comprises an area of 6,500 square miles with its topography varying greatly from the western sea-level plains to easterly mountainous terrain reaching heights of more than 13,000 feet.

The temperature in this province varies greatly by altitude, but maintains an average of 70-75° F year round. The climate also varies greatly by location—the windward sides of the islands experience heavy precipitation year round, while the leeward slopes are semiarid.

Hawaii is characterized by its tropical climate and volcanic history–all 9 islands are volcanoes at various stages of activity and decay. Due to its isolation and unique soil profile, the Hawaiian Islands Province has a small but distinctive population of fauna. There are many native bird species, along with a few species of introduced mammals. The flora includes many endemic species, and major habitat types include shrublands, forests, bogs, and moss and lichen covered mountainsides and outcrops that vary according to elevation and wind exposure.

Long before there were homes, development, and invasive species in this area, the original, native vegetation provided continuous cover and adjacent feeding opportunities for wildlife, including pollinators. In choosing plants, aim to create habitat for pollinators that allow adequate food, shelter, and water sources. Most pollinators have very small home ranges. You can make a difference by understanding the vegetation patterns of the farm, forest, or neighbor's yard adjacent to you and by making planting choices that support the pollinators' need for food and shelter as they move through the landscape.

UNDERSTANDING



Photo courtesy Colleen Kwong

THE HAWAIIAN ISLANDS PROVINCE

- This region is designated M423 in the Baileys' Ecosystem Provinces. To see a map of the provinces go to: www.fs.fed.us/rm/ecoregions/products/map-ecoregionsunited-states/.
- Not sure about which bioregion you live or work in? Go to www.pollinator.org and click on Ecoregion Locator for help.
- **%** The Hawaiian Islands region covers 6,500 square miles.
- Primarily a hilly and mountainous region, all the islands are volcanoes in various stages of activity and decay.
- **ﷺ** Elevations ranging from sea level to higher than 13,000 feet.
- **%** Average annual temperature ranges from 70° to 75° F.
- X Year-round precipitation can reach averages of 200 inches.
- 🕷 USDA Hardiness Zones 9a-13a (2012 version).

CHARACTERISTICS

- **%** The ohia tree, its accompanying ferns, and the koa tree predominate ecosystems unique to Hawaii's wet forests.
- Shrublands exist primarily on leeward mountainsides where rainfall is minimal.
- **ﷺ** Bogs are common at higher elevations with heavy rainfall.
- X Mosses and lichens survive even above the tree line, where frost is a frequent occurrence.
- X The volcanic islands' surfaces vary from deep Ultisols on the older islands to igneous rock on the youngest islands.



The Hawaiian Islands Province includes:

The Tropical Moist Forest, Tropical Dry Forest, and Tropical Grassland and Shrubland "CREATING POLLINATOR HABITAT IS ONE PRACTICE WE MUST EMPLOY TO MITIGATE THE IMPACT OF NEW VIRAL AND PARASITIC PRESSURES ON HAWAII'S ENVIRONMENTALLY ISOLATED FERAL AND DOMESTICATED HONEY BEE POPULATIONS."

-- DAVE SCHELL, GM, ISLAND PRINCESS MACADAMIA NUT COMPANY, KEA'AU, HAWAII, OPERATIONS

MEET THE POLLINATORS



I'iwii bird

Photo courtesy Karl Magnacca



WHO ARE THE POLLINATORS?

BEES

Bees are well documented pollinators in the agricultural and natural systems of the Hawaiian Islands Province. A wide range of crops including Macadamia nuts, watermelon, guava, and coffee all benefit from bee pollination.

Most of us are familiar with the colonies of honey bees that have been the workhorses of agricultural pollination for years in the United States. They were imported to Hawaii from Europe about 150 years ago.

There are nearly 4000 species of native ground and twig nesting bees in the U.S. Some form colonies while others live and work a solitary life. Native bees currently pollinate many crops and can be encouraged to do more to support agricultural endeavors if their needs for nesting habitat are met and if suitable sources of nectar, pollen, and water are provided. Bees have tongues of varying lengths that help determine which flowers they can obtain nectar and pollen from.

The **sweat bee** (family *Halictidae*) nests underground. Various species are solitary while others form loose colonies.

Solitary bees include carpenter bees (*Xylocopa* spp.), which nest in wood; digger, or polyester bees (*Colletes* spp.), which nest underground; leafcutter bees (*Megachile* spp.), which prefer dead trees or branches for their nest sites; and mason bees (*Osmia* spp.), which utilize cavities that they find in stems and dead wood.

BUTTERFLIES

Gardeners have been attracting butterflies to their gardens for some time. These insects tend to be evecatching, as are the flowers that attract them. Position flowering plants where they have full sun and are protected from the wind. Also, you will need to provide open areas (e.g. bare earth, large stones) where butterflies may bask, and moist soil from which they may get needed minerals. By providing a safe place to eat and nest, gardeners can also support the pollination role that butterflies play in the landscape. It might mean accepting slight damage to the plants, known as host plants, that provide food for the larval stage of the butterfly.

A diverse group of butterflies are present in garden areas and woodland edges that provide bright flowers, water sources, and specific host plants. Numerous trees, shrubs, and herbaceous plants support butterfly populations.

Butterflies are in the Order *Lepidoptera*. Some of the species in the Hawaiian Islands Province are the Citrus Swallowtail, Painted Lady, Hawaiian Blue, and the King Kamehameha butterfly. They usually look for flowers that provide a good landing platform.

Wet mud areas provide butterflies

with both the moisture and minerals they need to stay healthy. Butterflies eat rotten fruit and even dung, so don't clean up all the messes in your garden!

MOTHS

Moths are most easily distinguished from butterflies by their antennae. Butterfly antennae are simple with a swelling at the end. Moth antennae differ from simple to featherlike, but never have a swelling at the tip. In addition, butterflies typically are active during the day; moths at night. Butterfly bodies are not very hairy, while moth bodies are quite hairy and more stout.

Moths, generally less colorful than butterflies, also play a role in pollination. They are attracted to flowers that are strongly sweet smelling, open in late afternoon or night, and are typically white or pale colored.

BEETLES

Over 30,000 species of beetles are found in the United States and many of them can be found on flower heads. Gardeners have yet to intentionally draw beetles to their gardens, possibly because beetle watching isn't as inspiring as butterfly or bird watching. Yet beetles do play a role in pollination. Some have a bad reputation because they can leave a mess behind, damaging plant parts that they eat. Beetles are not as efficient as some pollinators. They wander between different species, often dropping pollen as they go. Beetle pollinated plants tend to be

large, strong scented flowers with their sexual organs exposed. They are known to pollinate Magnolia, sweetshrub (*Calycanthus*), paw paws, and yellow pond lilies.

FLIES

It may be hard to imagine why one would want to attract flies to the garden. However, like beetles, the number of fly species and the fact that flies are generalist pollinators (visit many species of plants), should encourage us all to leave those flies alone and let them do their job as pollinators.

Recent research indicates that flies primarily pollinate small flowers that bloom under shade and in seasonally moist habitats. The National Research Council's *Status of Pollinators in North America* study states that flies are economically important as pollinators for a range of annual and bulbous ornamental flowers.

Plants pollinated by the fly include the American pawpaw (Asimina triloba), dead horse arum (Helicodiceros muscivorus), skunk cabbage (Symplocarpus foetidus), mango (Mangifera indica), lychee (Litchi chinensis), and members of the carrot family like Queen Anne's lace (Daucus carota).

BIRDS

The I'iwi bird is known to pollinate the Ohia tree.Like many Hawaiian honeycreepers, the I'iwi bird is nectarviovous. These birds use their pointy bills to collect and feed on nectar, pollinating as they move from

plant to plant.

The brightly colored and aptly named Hummingbird Moth often causes confusion; but there are no hummingbirds, either native or introduced, in the Hawaiian Islands Province.

However, hummingbirds are the primary birds which play a role in pollination in the rest of North America. Their long beaks and tongues draw nectar from tubular flowers. Pollen is carried on both the beaks and feathers of different hummingbirds. The regions closer to the tropics, with warmer climates, boast the largest number of hummingbird species and the greatest number of native plants to support the bird's need for food. Whitewinged doves (Zenaida asiatica) are also pollinators of the saguaro cactus (Carnegeia gigantea) in the south central United States.

Bright colored tubular flowers attract hummingbirds to gardens throughout the United States. Hummingbirds can see the color red; bees cannot.

BATS

Though Hawaii's lone native bat, the Hawaiian Hoary Bat, is primarily an insectivore, bats play an important role in pollination in the mainland southwest where they feed on agave and cactus. The longnosed bats' head shape and long tongue allows it to delve into flower blossoms and extract both pollen and nectar.

WHICH FLOWERS DO THE POLLINATORS PREFER?

NOT ALL POLLINATORS ARE found in each North American province, and some are more important in different parts of the United States. Use this page as a resource to understand the plants and pollinators where you live.

Plants can be grouped together based on the similar characteristics of their flowers. These floral characteristics can be useful to predict the type of pollination method or animal that is most effective for that group of plants. This association between floral characteristics and pollination method is called a pollination syndrome.

The interactions of animal pollinators and plants have influenced the evolution of both groups of organisms. A mutualistic relationship between the pollinator and the plant species helps the pollinator find necessary pollen and nectar sources and helps the plant reproduce by ensuring that pollen is carried from one flower to another.

PLANT TRAITS

| Plant | | | | | |
|------------------|--|---|-------------------------------------|--|--|
| Trait | Bats | Bees | Beetles | | |
| Color | Dull white, green or purple | Bright white, yellow, blue, or UV | Dull white or green | | |
| Nectar guides | Absent | Present | Absent | | |
| Odor | Strong musty; emitted at night | Fresh, mild, pleasant | None to strongly fruity or fetid | | |
| Nectar | Abundant; somewhat hidden | Usually present | Sometimes present; not hidden | | |
| Pollen | Ample | Limited; often sticky and scented | Ample | | |
| Flower Shape | Regular; bowl shaped – closed during day | Shallow; have landing platform; tubular | Large bowl-like, Magnolia | | |

This chart and more information on pollinator syndromes can be found at:



AND THE POLLINATORS THEY ATTRACT

Pollinator

| Birds | Butterflies | Flies | Moths | Wind |
|---|---|--|---|--|
| Scarlet, orange, red or white | Bright, including red and purple | Pale and dull to dark brown or purple; flecked with translucent patches | Pale and dull red, purple, pink or white | Dull green, brown, or colorless; petals absent or reduced |
| Absent | Present | Absent | Absent | Absent |
| None | Faint but fresh | Putrid | Strong sweet; emitted at night | None |
| Ample; deeply hidden | Ample; deeply hidden | Usually absent | Ample; deeply hidden | None |
| Modest | Limited | Modest in amount | Limited | Abundant; small, smooth, and not sticky |
| Large funnel like; cups, strong perch support | Narrow tube with spur; wide landing pad | Shallow; funnel like or complex and trap-like | Regular; tubular without a lip | Regular; small and stigmas exerted |

http://www.fs.fed.us/wildflowers/pollinators/syndromes.shtml

DEVELOPING LANDSCAPE PLANTINGS THAT PROVIDE POLLINATOR HABITAT

WHETHER YOU ARE A FARMER of many acres, land manager of a large tract of land, or a gardener with a small lot, you can increase the number of pollinators in your area by making conscious choices to include plants that provide essential habitat for bees, butterflies, moths, beetles, and other pollinators.

FOOD:

Flowers provide nectar (high in sugar and necessary amino acids) and pollen (high in protein) to pollinators.

Fermenting fallen fruits also provide food for bees, beetles and butterflies. Specific plants, known as host plants, are eaten by the larvae of pollinators such as butterflies.

- Plant in groups to increase pollination efficiency. If a pollinator can visit the same type of flower over and over, it doesn't have to relearn how to enter the flower and can transfer pollen to the same species, instead of squandering the pollen on unreceptive flowers.
- Plant with bloom season in mind, providing food year round (see Bloom Periods pp. 16-17).
- Plant a diversity of plants to support a variety of pollinators. Flowers of different color, fragrance, and season of bloom on plants of different heights will attract different pollinator species and provide pollen and nectar throughout the seasons.

• Many herbs and annuals, although not native, are very good for pollinators. Mint, sage, basil, rosemary, oregano, lavender, garlic, and chives area just a few herbs that can be planted. Old fashioned zinnias, cosmos, and single sunflowers support bees and butterflies.

• Learn and utilize Integrated Pest Management (IPM) practices to address pest concerns. Minimize or eliminate the use of pesticides.

SHELTER:

Pollinators need protection from severe weather and from predators as well as sites for nesting and roosting.

- Incorporate different canopy layers in the landscape by planting trees, shrubs, and different-sized perennial plants.
- Leave dead snags for nesting sites of bees, and other dead plants and leaf litter for shelter.

• Build bee boxes to encourage solitary, non-aggressive bees to nest on your property.

• Leave some areas of soil uncovered to provide ground nesting pollinators easy access to underground tunnels.

• Group plantings so that pollinators can move safely through the landscape protected from predators.

• Include plants that are needed by butterflies during their larval development.

WATER:

A clean, reliable source of water is essential to pollinators.

- Natural and human-made water features such as running water, pools, ponds, and small containers of water provide drinking and bathing opportunities for pollinators.
- Ensure the water sources have a shallow or sloping side so the pollinators can easily approach the water without drowning.

Your current landscape probably includes many of these elements. Observe wildlife activity in your farm fields, woodlands, and gardens to determine what actions you can take to encourage other pollinators to feed and nest. Evaluate the placement of individual plants and water sources and use your knowledge of specific pollinator needs to guide your choice and placement of additional plants and other habitat elements. Minor changes by many individuals can positively impact the pollinator populations in your area. Watch for - and enjoy - the changes in your landscape!

• CAUTION: Remember that pesticides are largely toxic to pollinators. Extreme caution is warranted if you choose to use any pesticide. Strategically apply pesticides only for problematic target species. Read the label. Look for bee caution statements.



Macadamia nuts, watermelon, guava, and coffee are a few of the food crops in the Hawaiian Islands Province that will benefit from strong native bee populations that boost pollination efficiency. Incorporate different plants throughout the farm that provide food for native populations when targeted crops are not in flower.

Farmers have many opportunities to incorporate pollinator-friendly land management practices on their land which will benefit the farmer in achieving his or her production goals:

• Manage the use of pesticides to reduce the impact on native pollinators. Spray when bees aren't active (just after dawn) and choose targeted ingredients.

- Carefully consider the use of herbicides.
- Minimize tillage to protect ground nesting pollinators.
- Ensure water sources are scattered throughout the landscape.
- Choose a variety of native plants to act as windbreaks, riparian buffers, and field borders throughout the farm.
- Plant unused areas of the farm with temporary cover crops that can provide food or with a variety of trees, shrubs, and flowers that provide both food and shelter for pollinators.
- Check with your local Natural Resources Conservation Service (NRCS) office to see what technical and financial support might be available to assist you in your effort to provide nectar, pollen, and larval food sources for pollinators on your farm.



"ADDING NATIVE PLANTINGS TO IMPROVE POLLINATOR HABITAT MAKES SENSE IN ADVANCING OUR FAMILY FARM'S CONSERVATION AND ECONOMIC OBJECTIVES, ENHANCING BENEFICIAL WILDLIFE AND IMPROVING POLLINATION IN OUR ORCHARD AND "

--LEE MCDANIEL, FARMER AND PRESIDENT, NATIONAL ASSOCIATION OF CONSERVATION DISTRICTS

PUBLIC LANDS

HAWAII'S ISOLATION
HAS FOSTERED
UNIQUE ECOSYSTEMS.
POLLINATOR
CONSERVATION
SUPPORTS THESE
THREATENED
COMMUNITIES AND
IS ALSO INTEGRAL TO
FOOD SECURITY, AN
IMPORTANT ISLAND
CONSIDERATION."

-DANIELLE DOWNEY, APICULTURE SPECIALIST, HAWAII DEPARTMENT OF AGRICULTURE



Public lands are maintained for specific reasons ranging from high impact recreation to conservation. In the Hawaiian Islands Province many ecosystems have been destroyed or modified to allow for the expansion of development, especially within the tourism industry. This includes the alteration of watersheds and other natural areas to allow for roads, buildings, marinas, and man-made landscapes. Less disturbed natural areas can be augmented with plantings of native plant species. Existing plantings around buildings and parking areas should be evaluated to determine if pollinator-friendly plants can be substituted or added to attract and support pollinators. Public land managers have a unique opportunity to use their plantings as an education tool to help others understand the importance of

pollinators in the environment through signs, brochures, and public programs.

In an effort to increase populations of pollinators the land manager can:

- Inventory and become
- knowledgeable of local pollinators.

• Provide connectivity between vegetation areas by creating corridors of perennials, shrubs, and trees that provide pollinators shelter and food as they move through the landscape.

- Maintain a minimum of lawn areas that support recreational needs.
- Restrict the use of pesticides and herbicides.
- Provide water sources in large open areas.
- Maintain natural meadows and openings that provide habitats for sun-loving wildflowers and grasses.
- Remove invasive species and encroaching shrubs and trees.

HOME LANDSCAPES



"A GARDEN IS ONLY AS RICH AND BEAUTIFUL AS THE INTEGRAL HEALTH OF THE SYSTEM; POLLINATORS ARE ESSENTIAL TO THE SYSTEM - MAKE YOUR HOME THEIR HOME."

-- DERRY MACBRIDE NATIONAL AFFAIRS AND LEGISLATION CHAIRWOMAN, GARDEN CLUB OF AMERICA Gardeners have a wide array of plants to use in their gardens. Native plants, and plants developed by professional and amateur breeders can be found in garden centers, in catalogs, and on websites. Use your knowledge of pollinator needs to guide your choices.

- Choose a variety of plants that will provide nectar and pollen throughout the growing season.
- Resist the urge to have a totally manicured lawn and garden. Leave bare ground for ground nesting bees. Leave areas of dead wood and leaf litter for other insects.
- Strive to eliminate the use of all pesticides.
- Find local resources to help you in your efforts. Contact your local county extension agent or native plant society. Visit your regional botanic gardens and arboreta.



but it is important to remember that you are trying to provide connectivity to the landscape adjacent to your property. Don't just look within your property boundaries. If your neighbor's property provides an essential element, such as water, which can be utilized by pollinators visiting your land, you may be able to devote more space to habitat elements that are missing nearby. It is best to use native plants which have evolved to support the needs of specific native pollinators. Some pollinators, however, are generalists and visit many different plants, both native and non-native. Be sure that any non-native plants you choose to use are not invasive. Remember that specialized cultivars sometimes aren't used by pollinators. Flowers that have been drastically altered, such as those that are double or a completely different color than the wild species, often prevent pollinators from finding and feeding on the flowers. In addition, some altered plants don't contain the same nectar and pollen resources that attract pollinators to the wild types.

The scale of your plantings will vary

• CAUTION: Take time to evaluate the source of your plant material. You want to ensure you get plants that are healthy and correctly identified. Your local native plant society can help you make informed decisions when searching for plants.



PLANTS THAT ATTRACT POLLINATORS IN THE HAWAIIAN ISLANDS PROVINCE

The following chart lists plants that attract pollinators, are native to the Hawaiian Islands, and are available for purchase in the nursery trade. Not every plant listed is native to every Hawaiian Island, so try to select plants native to where they will be planted. This is not an exhaustive list, but provides guidance on where to start. Annuals, herbs, and cover crops provide food and shelter for pollinators, too. Unlike our other ecoregional planting guides, this particular guide includes plants that are rare, threatened, or endangered. Before purchasing plants, check for the official Department of Land and Natural resources tag that certifies that plants were collected and propagated legally.

| Scientific Name | Common Name | Habit | Flower Color | Height (in feet) | Perennation and Bloom Period | Sun | Soil | Elevation (in feet) | Pollinators | Also a host |
|--|------------------------------------|---------------|---------------------------------------|---------------------|---------------------------------------|-----------------------------|-----------------------------|---------------------------------------|---|----------------|
| | | | | Fo | orbs | | | | | |
| Argemone glauca | pua kala | Forb | White and Yellow | 2-6' | Perennial | Full Sun | Well Drained | 0-6200 | bees, wasp | |
| Astelia menziesiana | pua'akuhinia | Forb | Purple, Red, White, and Yellow | 1-6′ | Pernnial, April- June | Partial Sun - Shade | Well Drained | High wet forests only 2000-7000 | bees, flies | |
| Bacopa monnieri | water hyssop | Forb | Pink, Purple, and White | <1′ | Perennial, Year Round | Full Sun – Partial Shade | Moist | 0-300 | bees, flies | |
| Bidens hawaiensis | ko`oko`olau | Forb/Subshrub | Yellow | 2-6' | Perennial, Year Round | Full Sun | Moist | 1600-4600 | bees, flies (Bombilid bee flies) | |
| Bidens torta | koʻokoʻolau | Forb/Subshrub | Yellow | 4-10' | Perennial | Full Sun | Well Drained to Moist | 650-4000 | bees, flies (Bombilid bee flies) | |
| Dianella sandwicensis | 'uki'uki | Forb | Light Blue to White | 1-3′ | Perennial, Year Round | Full Sun – Partial Shade | Dry to Moist | 400-7000 | bees, beetles | |
| Heliotropium anomalum | Polynesian heliotrope | Forb/Subshrub | Light Purple to White | <2' | Perennial, Year Round | Full Sun | Well Drained | Coastal | bees (small carpenter), small butterflies (skippers) | |
| Heliotropium curassavicum | salt heliotrope | Forb/Subshrub | Light Purple/Light Yellow to White | <2' | Annual or Perennial, Year Round | Full Sun | Well Drained | <6800 | bees | |
| Lipochaeta integrifolia (Melanthera integrifolia) | nehe | Forb/Subshrub | Yellow | < 1′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | <150 | bee | |
| Lobelia hypoleuca | opelu | Forb | Blue-Purple | 6-12' | Perennial | Full Sun – Partial Shade | Dry-Moist | 2000-5000 | bee, moth (hummingbird moth), butterfly | |
| Phyllostegia macrophylla | Large leaved mint | Forb | White | 3 | Perennial | Shade | Well Drained - Moist | 2400-4000' | bees, flies | |
| Phytolacca sandwicensis | Popolo ku mai-Hawaiian pokeweed | Forb | Pink, White | 2-10+' | Perennial | Partial Sun | Well Drained- Wet, Moist | 2200-6500 | flies | |
| Portulaca lutea | native yellow purslane | Forb | Yellow | 1′ | Perennial, July- November | Full Sun | Well Drained | 0-130 | bees, flower flies | |
| Portulaca molokiniensis | ʻihi | Forb | Yellow | 1-1.5′ | Perennial, Year Round | Full Sun | Well Drained | 0-1000 | bees, flower flies | |
| Portulaca sclerocarpa | 'ihi makole | Forb | White - Pink | >1' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 3300-5300 | bees, flower flies | |
| Portulaca villosa | hairy purslane | Forb | White | 1-2.5′ | Perennial, Year Round | Full Sun | Well Drained | 0-2000 | bees, flower flies | |
| Scaevola coriacea | dwarf naupaka | Forb/Subshrub | White | <2' | Perennial, Year Round | Full Sun | Well Drained | 0-150 | bees, flies, moth (with a long tongue) | |
| Sesuvium portulacastrum | shoreline seapurslane | Forb | Pink | <1′ | Perennial, Year Round | Full Sun | Well Drained to Wet | 0-150 | bees, flies | |
| Waltheria indica | `uhaloa | Forb/Subshrub | Yellow | Up to 6' | Perennial | Full Sun | Well Drained to Moist | 0-4000 | bees, butterflies | |
| | | | | Vi | nes | | | | | |
| Bidens cosmoides | koʻokoʻolau | Vine/Forb | Yellow | Vine | Perennial, Year Round | Full Sun – Partial Sun | Moist | High elevation | Birds, flies (bee flies), bees | |

(continued on page 17)

| Scientific Name | Common Name | Habit | Flower Color | Height (in feet) | Perennation and Bloom Period | Sun | Soil | Elevation (in feet) | Pollinators | Also a host |
|-------------------------|---------------------------|----------------|------------------------|---------------------|---|--------------------------------|--------------------------|--------------------------|------------------------------------|---|
| Canavalia galeata | 'awikiwiki | Vine/Forb | Pink-Purple | Vine | Perennial | Partial Sun | Moist | 600-2600 | bees | |
| Canavalia hawaiiensis | puakauhi | Vine/Forb | Pink-Purple | Vine | Perennial | Full Sun – Partial Sun | Moist | 0-1000 | bees | |
| Canavalia kauaiensis | Kauai jackbean | Vine/Forb | Pink | Vine | Perennial | Filtered | Well Drained - Moist | 600 - 2600 | bees | |
| Canavalia pubescens | lavafield jackbean | Vine/Forb | Pink-Purple | Vine | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 0-2000 | bees | |
| Freycinetia arborea | `ie`ie | Vine/Forb | Orange | Vine | Perennial, Year Round | Full Sun – Partial Sun | Well Drained - Moist | 1000-5000 | Birds, bees, wasps | beetles will likely eat pollen |
| lpomoea pes-caprae | p huehue | Vine/Forb | Purple | Vine | Perennial, Year Round | Full Sun | Well Drained | 0-2000 | bees , flies | |
| Jacquemontia ovalifolia | pa'o Hi'iaka | Vine/Forb | Blue-White | Vine | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 0-100 | bees, flies | |
| Smilax melastomifolia | `hoi | Vine/Forb | White | Vine | Perennial | Shade | Well Drained - Wet | 600-6800 | bees, flies, moth | moth |
| Vigna marina | m hihihi | Vine/Forb | Yellow | Vine | Perennial, Sporatic | Full Sun – Partial Sun | Well Drained | 0-1000 | bees | |
| | | | | Trees an | d Shrubs | | | | | |
| Abutilon eremitopetalum | hiddenpetal Indian mallow | Shrub | Green-Red | 2'-10' | Perennial, February | Full Sun – Partial Sun | Well Drained | 700-1700 | bees, birds, butterflies | |
| Abutilon menziesii | ko'oloa 'ula | Shrub | Pink-Red/Yellow | 2'-10' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 650-1700 | birds, butterflies, bees | |
| Acacia koa | koa | Tree | Yellow | 15'- > 50' | Perennial, January-March | Full Sun - Partial Sun | Well Drained | 200-6500 | bees, flies | |
| Acacia koaia | koai'a | Tree | Yellow | 76'-96' | Perennial, January-March | Full Sun - Partial Sun | Well Drained | 150-4000 | bees, flies | |
| Alyxia stellata | maile | Shrub/Vine | Green-White/ Yellow | 2'-10' | Perennial, Fall, Winter | Full Sun – Partial Sun | Well Drained- Moist | 150-5000 | moths, bees | |
| Antidesma platyphyllum | Hame | Shrub/Tree | Green-White | 15'-50' | Perennial, June- October | Partial Sun | Well Drained- Moist | 1500-4000 | bees, flies | |
| Antidesma pulvinatum | hame | Tree | Cream | 15'-50' | Perennial, February-October | Full Sun – Partial Sun | Well Drained-Dry, Wet | 0-5000 | bees, flies | |
| Artemisia mauiensis | Maui wormwood | Shrub/Subshrub | Brown | 2'-6' | Perennial | Full Sun – Partial Sun | Well Drained | 6200-7500 | bees, flies, butterflies, moths | |
| Brighamia insignis | 'ālula | Shrub | Yellow | 2'-15' | Perennial, September- October | Full Sun – Partial Sun | Well Drained | 0-1300 | moths, bees | |
| Broussaisia arguta | kanawao | Shrub/Tree | Cream-Blue-Pink | 6'-19' | Perennial | Partial Sun | Well Drained- Moist | 1000-6000 | bees, flies, wasps, moth | |
| Caesalpinia kavaiensis | ʻuhi'uhi | Tree | Pink-Red | 10'-30' | Perennial, Year Round | Full Sun | Well Drained | 250-3000 | bees, moths | |
| Capparis sandwichiana | maiapilo | Shrub | White | 2'-10' | Perennial, Spring- Summer | Full Sun | Well Drained | 0-325 | moths, bees | |
| Cheirodendron trigynum | ʻōlapa | Tree | Purple | 19'-57' | Perennial | Shade | Well Drained to Wet | 1600-7000 | birds, bees | |
| Claoxylon sandwicense | po'ola | Shrub | White- Pale Green | up to 8.5' | Perennial | Partial Sun – Partial Shade | Well Drained | Dry and mesic forests | bees, birds | |
| Clermontia hawaiiensis | `ohawai nui | Shrub | Green-White | 2'-10' | Perennial, Feburary-March and June-July | Partial Sun – Partial Shade | Moist | 1000-5000 | bees, birds | beetles will likely eat pollen |
| Clermontia montis-loa | `ohawai | Shrub | Pink-Purple | 6-15' | Perennial, Year Round | Full Sun – Partial Shade | Well Drained - Moist | 2500-6000 | bees, birds | beetles will likely eat pollen |
| Clermontia parviflora | `ohawai | Shrub | Green-Purple- White | 15'-30' | Perennial | Partial Sun | Well Drained- Moist | 400-4600 | bees, birds | beetles will likely eat pollen |

(continued on page 18)

(continued from page 17) PLANTS THAT ATTRACT POLLINATORS IN THE HAWAIIAN ISLANDS PROVINCE

| Scientific Name | Common Name | Habit | Flower Color | Height (in feet) | Perennation and Bloom Period | Sun | Soil | Elevation (in feet) | Pollinators | Also a host |
|---|------------------------|-----------------|------------------------------|---------------------|------------------------------------|----------------------------|-------------------------------|------------------------|---|----------------|
| Coprosma ernodeoides | 'aiakanene | Shrub | Cream-Yellow | 2'-6' | Perennial, Spring-Summer | Full Sun – Partial Sun | Well Drained | 4000-8500 | wind pollinated (bees and moths will visit) | |
| Coprosma pubens | pilo | Tree | White | 10-30' | Perennial | Full Sun – Filtered Sun | Well Drained - Wet | 1200-6200 | wind pollinated (bees and moths will visit) | |
| Cyrtandra platyphylla | `ilihia | Shrub | White | 3′ | Perennial | Shade | Well Drained - Moist | 1200-4500 | flies | |
| Delissea rhytidosperma | Kauai delissea | Shrub | Green-White | 2-10′ | Perennial, June-December | Partial Sun | Well Drained- Moist | 980-3300 | bird, moth, bee | |
| Diospyros hillebrandii | elama | Tree | Cream | 15-50' | Perennial | Partial Sun – Shade | Well Drained | 490-2500 | bee, fly | |
| Diospyros sandwicensis | lama | Tree | White | 6-35′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 0-4000 | bee, fly | |
| Erythrina sandwicensis | wili wili | Tree | Orange-White- Yellow | 15-50′ | Perennial, Spring-Summer | Full Sun | Well Drained | 0-1950 | bird, butterfly, moth, bee | |
| Euphorbia celastroides var. kaenana | 'akoko | Shrub/Tree | Brown/White | 2-6′ | Perennial, Year Round | Full Sun | Well Drained | 0-1000 | bee, fly, moth | |
| Gardenia brighamii | nā'ū | Shrub/Tree | White | 6-30' | Perennial, Year Round | Full Sun | Well Drained | 1100-1700 | moth, bee | |
| Gossypium tomentosum | ma'o | Shrub | Yellow | 2-6′ | Perennial, Year Round | Full Sun | Well Drained | <400 | bee, moth, butterfly | |
| Hibiscus arnottianus | Oʻahu kokiʻo ke'oke'o | Shrub/Tree | White | 10-50' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 390-2500 | bird, moth, bee | |
| Hibiscus brackenridgei | ma'o hau hele | Shrub/Tree | Yellow | 2-10′ | Perennial, February-May | Full Sun – Partial Sun | Well Drained | 400-2600 | bird, moth, bee | |
| Hibiscus clayi | koki'o 'ula | Shrub/Tree | Red | 10-30' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 150-2000 | bird, moth, bee | |
| Hibiscus kokio | koki'o | Shrub/Tree | Red, Orange, Yellow | 2-50' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 500-3000 | bird, moth, bee | |
| Hibiscus waimeae | Kaua'i koki'o ke'oke'o | Shrub/Tree | White | 10-50' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 2000-3000 | bird, moth, bee | |
| llex anomala | kawa'u | Shrub/Tree | White | 10-50' | Perennial | Full Sun – Partial Sun | Well Drained- Moist | 2000-4500 | bee, fly | |
| Kadua affinis | manono | Tree/Shrub/Vine | Green, Purple | 6-30' | Perennial | Partial Sun | Well Drained- Moist | 850-6690 | moth | |
| Metrosideros polymorpha | 'ohi'a lehua | Shrub/Tree | Red, Pink, Orange, Yellow | 2-50+' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained Dry and Moist | 0 - 7200+ | bird, wasp, bee | |
| Myoporum sandwicense | naio | Shrub/Tree | White-Pink | 2-50+' | Perennial, Year Round | Full Sun | Well Drained | 0 - 7700+ | fly, bee | |
| Myrsine lessertiana | kolea lau nui | Shrub/Tree | Pink-Purple | 15-30′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 700-7200+ | moth, fly, bee | |
| Myrsine sandwicensis | kokea lau li'i | Shrub/Tree | Pink | 13-25′ | Perennial | Full Sun – Partial Sun | Well Drained - Moist/Wet | 1000-4800 | fly, moth, bee | |
| Nesoluma polynesicum (Sideroxylon polynesicum) | keahi | Shrub/Tree | Green-White | 6-50' | Perennial, May-September | Full Sun – Partial Sun | Well Drained | 425-2100 | fly, moth, bee | |
| Nestegis sandwicensis | olopua | Shrub/Tree | White, Yellow | 30-50+' | Perennial | Full Sun – Partial Sun | Well Drained | 100-4265 | bee, fly | |
| Ochrosia compta | holei | Tree | White | 15-50′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 975-1625 | moth | |
| Osteomeles anthyllidifolia | 'ū lei | Shrub | White | 2-30′ | Perennial, Spring, Winter | Full Sun – Partial Sun | Well Drained | 0-7600 | bees, flies | |
| Perrottetia sandwicensis | olomea | Shrub/Tree | Red | 6-50' | Perennial, Year Round | Partial Sun – Shade | Well Drained- Wet | 1000-4000 | flies | |
| Pisonia sandwicensis | aulu | Tree | Brown-White | 30-50' | Perennial, June, July, August | Full Sun – Partial Sun | Well Drained | 850-3400+ | bees, flies | |
| Pisonia umbellifera | pāpala kēpau | Shrub/Tree | White, Pink | 10-50' | Perennial | Full Sun – Partial Sun | Well Drained | 295-2625 | moth, fly, bee | |
| Pittosporum confertiflorum | ho'awa | Shrub/Tree | White, Cream | 2-50′ | Perennial | Full Sun – Partial Sun | Well Drained | 550-7000 | moth, fly, bee | |

(continued on page 19)

| | U 2/ | | | | | | | | | |
|----------------------------|-------------------------|----------------|-------------------------------------|---------------------|--|---------------------------|-----------------------------|------------------------|------------------------------------|----------------|
| Scientific Name | Common Name | Habit | Flower Color | Height (in feet) | Perennation and Bloom Period | Sun | Soil | Elevation (in feet) | Pollinators | Also a host |
| Pittosporum glabrum | ho'awa | Tree | White, Cream | 15-30′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 785-6465 | moth, fly, bee | |
| Pittosporum hawaiiense | ho'awa | Tree | Cream | 10-25′ | Perennial | Full Sun – Partial Sun | Well Drained- Moist/Wet | 1700-6000 | moth, fly, bee | |
| Pittosporum hosmeri | ho'awa | Tree | White, Cream | 10-30′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 1200-3500 | moth, fly, bee | |
| Polyscias hawaiensis | ʻohe'ohe | Tree | Red | 15-50+' | Perennial | Full Sun – Partial Sun | Well Drained- Moist | 500-3500 | fly, moth | |
| Polyscias racemosa | false 'ohe | Tree | Red, Yellow | 15-30′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 395-1300+ | fly, moth | |
| Polyscias sandwicensis | 'ohe makai | Tree | Purple | 15-50+' | Perennial, Fall | Full Sun – Partial Sun | Well Drained | 100-2600+ | fly, moth | |
| Pritchardia spp. | loulu | Tree | Yellow | 15-60' | Perennial | Full Sun | Well Drained | Variable | fly | |
| Psychotria greenwelliae | kõpiko nui | Tree | White | 15′ | Perennial | Full Sun – Partial Sun | Well Drained - Moist | 2000 - 4000 | fly, bee, moth | |
| Psychotria mariniana | kōpiko | Tree | White | 15-50+' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 200-4000 | moth, bee, fly | |
| Psychotria mauiensis | 'opiko | Shrub/Tree | White-Green | 10-45′ | Perennial | Partial Sun | Moist | 500-5000 | fly, bee, moth | |
| Psydrax odorata | alahe'e | Shrub/Tree | White | 10-50′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | 0-2700 | moth, bee, fly | |
| Rhus sandwicensis | neneleau | Tree | White | 6-30′ | Perennial | Full Sun – Partial Sun | Well Drained- Moist | 500-2500 | bees, flies, honey bee favorite | |
| Rubus hawaiensis | ʻakala | Subshrub | Pink | 6-10′ | Perennial, April- July | Full Sun – Partial Sun | Moist | 2000-10000 | bees | |
| Rumex giganteus | wet forest dock Pawale | Subshrub/Shrub | Pink-Red | Up to 6.5' | Perennial | Full Sun | Mesic-Wet | 2000-10000 | bee, fly, butterfly | |
| Rumex skottsbergii | pawale | Subshrub/Shrub | Cream-Green | 3-5' | Perennial | Full Sun | Well Drained | Variable | bee, fly, moth | |
| Santalum paniculatum | dry-mesic `iliahi | Shrub/Tree | Green to Yellow and Orange/Brown | 10-35′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 1500-6500 | bee, fly | |
| Santalum freycinetianum | ʻiliahi | Shrub/Tree | Red, Pink | 2-50′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 800-3000 | butterfly, bee, bird | |
| Sapindus oahuensis | lonomea | Shrub/Tree | White | 30-50'+ | Perennial, Spring, Summer and Winter | Full Sun – Partial Sun | Well Drained | 200-2000 | bee, fly, moth | |
| Scaevola chamissoniana | mountain naupaka (Puna) | Shrub/Tree | White | 2-10′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 1500-3500 | fly, bee, moth | |
| Scaevola gaudichaudiana | naupaka kuahiwi | Shrub | White | 2-30' | Perennial, Year Round | Partial Sun-Shade | Well Drained- Moist | 555-2600 | fly, bee, moth | |
| Scaevola kilaueae | huahekili uka | Shrub | White | 2-10' | Perennial | Full Sun | Well Drained | 3000-5000 | fly, bee, moth | |
| Scaevola sericea | beach naupaka | Shrub | White | 2-10' | Perennial, Year Round | Full Sun | Well Drained | Coastal | fly, bee, moth | |
| Sesbania tomentosa | ʻohai | Shrub | Orange, Red, Yellow | 2-30' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained | < 2500 | bird, bee | |
| Sophora chrysophylla | mamani | Shrub/Tree | Yellow | 6-50' | Perennial, July- November | Full Sun – Partial Sun | Well Drained | 1400-10000+ | bird, bee | |
| Styphelia tameiameiae | pūkiawe | Shrub | Yellow | Up to 15' | Perennial | Full Sun | Well Drained to Wet | 0-10000 | bees, flies | |
| Vaccinium calycinum | ohelo kau la'au | Shrub | Pink | 2-30' | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Wet, Moist | 1600-5800 | bees, flies | |
| Vaccinium reticulatum | ohelo 'ai | Shrub | Red | 2-10′ | Perennial, Year Round | Full Sun – Partial Sun | Well Drained- Moist | 2100-12000 | bees, flies | |
| Vitex rotundifolia | põhinahina | Shrub | Blue, Purple | 2-10′ | Perennial, Year Round | Full Sun | Well Drained | 0-50 | bee, butterfly | |
| Wikstroemia sandwicensis | Mesic Tree Akia | Shrub/Tree | Green, Yellow | 2-10′ | Perennial, Fall- Winter | Filtered Sun | Well Drained - Wet | 10 - 5000 | butterfly, moth, bee | |
| Wikstroemia phillyreifolia | Hawai'i 'ākia | Shrub/Tree | Yellow | 2-15′ | Perennial | Full Sun | Well Drained | 0-7500+ | butterfly, moth, bee | |
| Wikstroemia uva-ursi | hillside false ohelo | Shrub | Yellow | 2-6′ | Perennial, Fall- Winter | Full Sun – Partial Sun | Well Drained | 0-1375+ | moth, butterfly | |

BEE-POLLINATED PLANTS of the HAWAIIAN ISLANDS PROVINCE

| Common Bee-Pollinated Ornamentals and Crops | | | | | | Common Bee | -Pollinate | d Orname | entals and | Cr |
|---|--------------------|--------------------|------------|------------------|--|----------------|--------------------|--------------------|------------|---------|
| | Large Carpenter | Small Carpenter | Leafcutter | Yellow- faced | | | Large Carpenter | Small Carpenter | Leafcutter | Yo f |
| | FLOWERS | | | | | | C | ROPS | | |
| Aster | х | х | х | х | | Eggplant | х | | | |
| Beebalm | | | х | | | Cucumber | х | | | |
| Black-eyed Susan | х | | х | | | Squash | х | | | |
| Catnip | | х | | х | | Pumpkins | х | | | |
| Coneflower | х | | х | | | Gourds | х | | | |
| Cow parsley | | | | х | | Tomatoes | х | | | |
| Goldenrod | | x | х | х | | Thyme | | | х | |
| Irises | х | | | | | | TROPIC | AL FRUITS | | |
| Lavender | х | | х | | | Citrus | | х | х | |
| Monring glory | | х | | | | Cherimoya | | х | | |
| Oregano | | х | | х | | Durian | х | | | |
| Passion flowers | х | | х | | | Eucalyptus | х | х | х | |
| Penstemon | х | | х | | | Guava | | х | | |
| Phacelia | | x | х | | | Loquat | | х | х | |
| Potentilla | | | | х | | Lychee | | х | х | |
| Rose | х | x | | х | | Macadamia nut | | | х | |
| Rosemary | х | | х | х | | Mango | | х | х | |
| Sage | х | | х | | | Mangosteen | | х | | |
| Salvia | х | | х | | | Palms | | х | | |
| Saxifrage | | | | х | | Passion fruit | х | | х | |
| Sorrel | | x | | | | Rambutan | | х | | |
| Sunflowers | х | х | х | | | Soursop | | х | | |
| Verbena | | | х | | | Starfruit | | Х | | |
| Yarrow | | Х | х | х | | Surinam cherry | | | х | |
| | | | | | | | | | | |

Wild honey bee on bidens sp. courtesy Kim Dillman



"CONSERVE HAWAII'S UNIQUE NATIVE PLANTS BY INCLUDING THEM IN HOME LANDSCAPING. NATIVE PLANTS ADD BEAUTY, VARIETY AND FOOD SOURCES FOR NATIVE POLLINATORS AND HONEY BEES."

– KIM DILLMAN, BIG ISLAND PLANTS, VOLCANO, HAWAII

rops Yellowfaced

> X X

Х

Х

Х

Х



HAWAII NURSERIES THAT SPECIALIZE IN FORESTRY TREES, NATIVE OR INTRODUCED

Note that most have minimum orders of \$100 to \$500 and most grow to order.

State Tree Nursery Division of Forestry and Wildlife Kamuela, HI tel: (Hilo) (808) 974-4221

David and Victoria Antova H and S Farms (Ferns only) hsfarms@besthawaiianflowers.com 808-969-6364

Allie Atkins Lehua Lena Nursery PO Box 1479 Kea'au, HI 96749 tel/fax (808) 966-7975 alizon@hawaii.edu

Rick Barboza Hui Ku - Maoli Ola Native Hawaiian Plant Specialists 46-403 Haiku Rd. Kaneohe, HI 96744 (808) 295-7777 nativehawaiianplants@gmail.com www.hawaiiannativeplants.com

Mauna Ikena

Linda Burnham Larish P.O. Box 1337 Kea'au, HI 96749 maunaikena@gmail.com (808) 966-6337 Gordon Czernick Makani Gardens 1625 West Kuiaha Rd Haiku, HI 96708 tel: 808-572-6337 fax: 808-572-6337 email: koa234@yahoo.com

David DeEsch Hamakoa Nursery Koa and 'ōhi'a seedlings PO Box 1428 Honokaa, HI 96727 tel: 808-775-1663 egznhoney@lycos.com http://hamakoa.com/

Nick Koch Nursery Solutions Inc. PO Box 474 Paauilo HI 96776 (808) 776-9900 info@nurserysolutionsinc.com jr@hawaiiforest.com http://www.nurserysolutionsinc.com

Betsy Garties and Victoria Paul Aikane Nursery PO Box 459 Kapaau, HI 96755 info@aikanenursery.com 808-889-5906 Nina Gutmanis and Louis Bauguess Grow Native Corporation Box 753 Mt. View, HI 96771 tel. (808) 968-8350 fax (808) 968-6853 email: endflorahi@aol.com

Brian Kiyabu

Amy Greenwell Garden Native and Polynesian-introduced plants only; min. order \$500.00 ph: 808-323-3318 fax: 808-323-2394 brian.kiyabu@bishopmuseum.org http://www.bishopmuseum.org/ exhibits/greenwell/plantsales.html PO Box 1053 Captain Cook, HI 96704

Lelan Nishek Kaua'i Nursery & Landscaping 3-1550 Kaumuali'i Hwy. Lihue, HI 96766 tel. 808-245-7747 email: knl@kauainursery.com www.kauainursery.com

Anna Palomino Ho'olawa farms (natives only) Haiku, HI 96708 (808) 575-5099 apalomino@hawaii.rr.com http://www.hoolawafarms.com Ethan Romanchak Jonathan Keyser Native Nursery LLC (808) 878-8276 fax (808) 876-1020 email: info@mauinativenursery.com mailing address: PO Box 806 Kula, HI 96790 Street address: 1267 Naalae Rd Kula Maui HI tel. (808) 281-8494 (Ethan Romanchak) tel. (808) 250-3341 (Jonathan Keyser) www.mauinativenursery.com

Kerin Rosenberger Keep it Native Kaua'i, HI (808) 742-9894 kerinr@hawaiiantel.net (native plants only)

Jill Wagner Future Forests Nursery P.O. Box 847 Kailua Kona, HI 96745 Phone: 808-325-2377 email: trees@forestnursery.com www.forestnursery.com

Aileen Yeh Aileen's Nursery 942 W Kawailani St. Hilo, HI 96720 tel and fax: 808-959-8040 cell 808-936-2671 email: ayeh@hawaii.rr.com

HABITAT AND NESTING REQUIREMENTS OF NATIVE HAWAIIAN BEES

Large carpenter bees:

Soft dead wood, poplar, cottonwood or willow trunks and limbs, structural timbers including redwood. Depending on the species, there may be one or two brood cycles per year. These bees can be active all day even in the hottest weather.

Small carpenter bees:

Pithy stems including roses and blackberry canes. These bees are more active in the morning but can be found at other times.

Leafcutter bees:

Pre-existing circular tunnels of various diameters in dead but sound wood created by emerging beetles, some nest in the ground. Leave dead limbs and trees to support not just pollinators but other wildlife. Leafcutter bees can be seen foraging throughout the day even in hot weather.

Yellow-faced bees:

In dead stems. These bees are more active during morning hours.

A BASIC CHECKLIST

BECOME FAMILIAR WITH POLLINATORS IN YOUR LANDSCAPE.

- 🕷 Watch for activity throughout the day and the seasons.
- Keep a simple notebook of when and what comes to your garden. NOTE: It is not necessary to identify each species when you first get started. Simply note if it is a bee that likes the yellow flower that blooms in the fall.
- Consult a local field guide or web site (www.pollinator.org) when you are ready to learn more details.

ADD NATIVE PLANTS TO ATTRACT MORE NATIVE POLLINATORS.

- 🕷 List the plants you currently have in your landscape.
- Determine when you need additional flowers to provide nectar and pollen throughout the growing season.
- X Add plants that provide additional seasons of bloom, create variable heights for shelter, and attract the types of pollinators you want.
- Con't forget to include host plants that provide food and shelter for larval development.
- Contact your local native plant society or extension agent for more help.

USE POLLINATOR FRIENDLY LANDSCAPE PRACTICES TO SUPPORT THE POLLINATORS YOU ATTRACT.

- **W** Use Integrated Pest Management Practices to address pest concerns.
- Tolerate a little mess leave dead snags and leaf litter, keep areas bare for ground nesting pollinators, and leave some weeds that provide food for pollinators.
- **%** Provide safe access to clean water.

NOTICE THE CHANGES THAT YOU HAVE HELPED TO CREATE!

"MONARCH

BUTTERFLIES

NEVER FAIL TO CATCH

THE VISITOR'S EYE

AND ALWAYS LEAD

TO A TEACHABLE

MOMENT."

-- LOGAN LEE, PRAIRIE SUPERVISOR MIDEWIN NATIONAL TALLGRASS PRAIRIE



Many books, websites, and people were consulted to gather information for this guide. Use this list as a starting point to learn more about pollinators and plants in your area.

BAILEY'S ECOREGION MAPS

USDA Forest Service http://www.fs.fed.us/land/ ecosysmgmt/ecoreg1_home.html

POLLINATION/POLLINATORS

Pollinator Partnership www.pollinator.org

Natural Resources Conservation Service www.nrcs.usda.gov

North American Pollinator Protection Campaign www.nappc.org

USDA Forest Service www.fs.fed.us/wildflowers/pollinators/

Wild Farm Alliance www.wildfarmalliance.org

The Xerces Society www.xerces.org

Illinois Natural History Survey www.inhs.uiuc.edu

Buchmann, S.L. and G.P. Nabhan. 1997. *The Forgotten Pollinators* Island Press: Washington, DC.

Committee on the Status of Pollinators in North America. 2007. *Status of Pollinators in North America* The National Academies Press: Washington, DC.

NATIVE PLANTS

Plant Pono www.plantpono.org/

Plant Conservation Alliance www.nps.gov/plants

Lady Bird Johnson Wildflower Center www.wildflower.org/plants/

USDA Hardiness Zone Map www.usna.usda/Hardzone/

USDA, NRCS. 2007. The PLANTS Database www.plants.usda.gov, 19 July, 2007 National Plant Data Center, Baton Rouge, LA 70874-4490 USA

NATIVE BEES

National Sustainable Information Service

"Alternative Pollinators: Native Bees" by Lane Greer, NCAT Agriculture Specialist, Published 1999, ATTRA Publication #IP126 www.attra.ncat.org/attra-pub/ nativebee.html

Agriculture Research Service

Plants Attractive to Native Bees table www.ars.usda.gov/Research/docs. htm?docid=12052

BUTTERFLIES AND MOTHS

Opler, Paul A., Harry Pavulaan, Ray E. Stanford, Michael Pogue, coordinators. 2006. Butterflies and Moths of North America. Bozeman, MT: NBII Mountain Prairie Information Node. www.butterfliesandmoths.org/ (Version 07192007)

Pyle, Robert Michael. 1981. National Audubon Society Field Guide to Butterflies. Alfred A. Knopf: New York, NY.

North American Buterfly Association www.naba.org

FEEDBACK

We need your help to create better guides for other parts of North America. Please e-mail your input to **feedback@pollinator.org** or fax to **415-362-3070**.

- How will you use this guide?
- Do you find the directions clear? If not, please tell us what is unclear.
- Is there any information you feel is missing from the guide?
- **%** Any other comments?

THANK YOU FOR TAKING THE TIME TO HELP!

POLLINATOR PARTNERSHIP



















RESEARCH AND WRITING:

ELIZABETH L. LEY STEPHEN BUCHMANN, PH.D. GARY KAUFFMAN OLIVIA JACOBSEN MARY RAGER LINDSAY KWONG VICTORIA WOJCIK, PH.D.

EDITORIAL:

LAURIE DAVIES ADAMS MARY RAGER JENNIFER TSANG

PRODUCTION SUPERVISION:

MARY RAGER JENNIFER TSANG

DESIGN:

MARGUERITE MEYER

CONCEPT REVIEW:

Big Island Plants, Kim Dillman Bishop Museum, Clyde Imada Botanic Gardens Conservation International – US, Abby Hird Island Princess, Dave Schell Hawaii Department of Agriculture, Pat Conant, Danielle Downey, Jacqueline Robson, Lauren Rusert Hawaii Department of Land and Natural Resources, Lyman Perry Hawaii Invasive Species Council, Josh Atwood, Chuck Chimera, Patricia Clifford Starr Environmental, Forest and Kim Starr The University of Hawaii, Karl Magnacca

PHOTO CONTRIBUTORS:

Karl Magnacca Kim Dillman Lauren Rusert Danielle Downey Jonathan Starr Colleen Kwong

ORIGINAL CONCEPT CREATOR: Larry Stritch, Ph.D.

For a copy of this brochure, or for another region, visit www.pollinator.org

The Pollinator Partnership / North American Pollinator Protection Campaign

423 Washington St., 5th Floor, San Francisco, CA 94111 – 415-362-1137 www.pollinator.org 🄌 www.nappc.org