

The following list of resources includes peer reviewed publications, government, university, corporate, and NGO commissioned reports, and symposium proceedings discussing the impacts of neonicotinoid and similar pesticides to bees. This list is robust, but not exhaustive, if you have additional resources to add to this list please contact Vicki Wojcik at vw@pollinator.org.

P2 is presenting this list as an aggregated resource. We do not make any claims, either in favor or against, the research or opinions presented by the authors. Popular media, magazine, blog, and non-reviewed formats have been omitted from this list.

Updated December 12, 2013

Peer Reviewed Publications

Abramson, CI, Sokolowski, MBC, Brown, EA, & Pilard, S (2011) The effect of pymetrozine (Plenum WG-50®) on proboscis extension conditioning in honey bees (*Apis mellifera*: Hybrid var. Buckfast). *Ecotoxicology and environmental safety*, 1–9. doi:10.1016/j.ecoenv.2011.11.038

Adgaba, N, Al-ghamdi, A.A, Chernet, MH, Ali, YA, Ansari, MJ, Radloff, SE, & Howard, RH (2012) An experiment on comb orientation by Honey Bees (Hymenoptera: Apidae) in traditional hives. *Journal of Economic Entomology*, 105(3), 777–782.

Alaux, C, Brunet, JL., Dussaubat, C, Mondet, F, Tchamitchan, S, Cousin, M, Brillard, J, et al. (2010). Interactions between *Nosema* microspores and a neonicotinoid weaken honeybees (*Apis mellifera*). *Environmental microbiology*, 12(3), 774–82. doi:10.1111/j.1462-2920.2009.02123.x

Albajes R, López C, Pons X (2003) Predatory fauna in cornfields and response to imidacloprid seed treatment. *J Econ Entomol.* 96(6):1805-13.

Aliouane Y, El Hassani Athiamethoxam K, Gary V, Armengaud C, Lambin M, Gauthier M, (2009) Subchronic exposure of honey bees to sublethal doses of pesticides: effects on behavior. *Environ. Toxicol. Chem.* 28(1): 113–122. <http://cognition.ups-tlse.fr/productscientific/documents/papers/Aliouane%20ET&C%2008.pdf>

Aufauvre, J, Biron, DG, Vidau, C, Fontbonne, R, Roudel, M, Diogon, M, Viguès, B, et al. (2012) Parasite-insecticide interactions: a case study of *Nosema ceranae* and *fipronil* synergy on honeybee. *Scientific reports*, 2, 326. doi:10.1038/srep00326

Babendreier D, Kalberer N, Romeis J, Fluri P, & FBigler (2004) Pollen consumption in honey bee larvae: a step forward in the risk assessment of transgenic plants. *Apidologie* 35 (3) 293-300 DOI: 10.1051/apido:2004016

Badiou-Beneteau, Alexandra; Carvalho, Stephan M.; Brunet, Jean-Luc; Carvalho, Geraldo A.; Bulete, Audrey; Giroud, Barbara; Belzunces, Luc P. (2012) Development of biomarkers of exposure to xenobiotics in the honey bee *Apis mellifera*: Application to the systemic insecticide thiamethoxam. *ECOTOXICOLOGY AND ENVIRONMENTAL SAFETY* 82(22-31):10.1016/j.ecoenv.2012.05.005

Bahlai, CA, et al (2010) Choosing organic pesticides over synthetic pesticides may not effectively mitigate environmental risk in soybeans. <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0011250>

Belzunces LP (1995) *A mellifera* L exposed to sublethal does of Deltamethrin. *Environmental Toxicology*, 14(5), 855–860.

Bernal, J., E. Garrido-Bailón, et al. (2010). "Overview of Pesticide Residues in Stored Pollen and Their Potential Effect on Bee Colony (*Apis mellifera*) Losses in Spain." *Journal of Economic Entomology* 103(6): 1964-1971.

Biddinger, DJ; Robertson, JL; Mullin, C; Frazier, J; Ashcraft, SA; Rajotte, EG; Joshi, NK; Vaughn, M (2013) Comparative Toxicities and Synergism of Apple Orchard Pesticides to *Apis mellifera* (L.) and *Osmia cornifrons* (Radoszkowski) *PLOS ONE* 8(9):e72587; 10.1371/journal.pone.0072587

- Blacquièrè, T, Smagghe G, van Geste CAM, & Mommaerts V (2012). Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology* (London, England). doi:10.1007/s10646-012-0863-x
- Boily, M; Sarrasin, B; DeBlois, C; Aras, P; Chagnon, M (2013) Acetylcholinesterase in honey bees (*Apis mellifera*) exposed to neonicotinoids, atrazine and glyphosate: laboratory and field experiments. *ENVIRONMENTAL SCIENCE AND POLLUTION RESEARCH* 20(8):5603-5614; 10.1007/s11356-013-1568-2
- Boncristiani H, Underwood R, Schwarz R, Evans JD, Pettis JS, & Vanengelsdorp D (2011) Direct effect of acaricides on pathogen loads and gene expression levels of honey bee *Apis mellifera*. *Journal of insect physiology*, (December). doi:10.1016/j.jinsphys.2011.12.011
- Bonmatin JM, Marchand PA, Charvet R, Moineau I, Bengsch ER, Colin ME (2005) Quantification of imidacloprid uptake in maize crops. *J Agr Food Chem* 53: 5336–5341.
- Bonmatin JM, Moineau I, Charvet R, Fleche C, Colin ME and ER Bengsch (2003) A LC/APCI-MS/MS Method for Analysis of Imidacloprid in Soils, in Plants, and in Pollens *Anal. Chem.* 75, 2027-2033
- Bonmatin, M, Moineau I, Charvet, R. Colin ME, Fleche C & ER Bengsch (2005) Behaviour of Imidacloprid in Fields. Toxicity for Honey Bees. In *Environmental Chemistry, Green Chemistry and Pollutants in Ecosystems*, Lichtfouse F, Schwarzbauer J, and Robert D (eds), New York: Springer pp 483-494
- Bonzini S, Tremolada P, Bernardinelli I, Colombo M, & Vighi M (2011) Predicting pesticide fate in the hive (part 1) experimentally determined τ -fluvalinate residues in bees, honey and wax. *Apidologie*, 42, 378–390.
- Boutabet K., Kebsa W, Alyane M, & Lahouel M (2011) Polyphenolic fraction of Algerian propolis protects rat kidney against acute oxidative stress induced by doxorubicin. *Indian journal of nephrology*, 21(2), 101–6. doi:10.4103/0971-4065.82131
- Brittain, C. a., Vighi, M., Bommarco, R., Settele, J., & Potts, S. G. (2010) Impacts of a pesticide on pollinator species richness at different spatial scales. *Basic and Applied Ecology*, 11(2), 106–115. doi:10.1016/j.baae.2009.11.007
- Brittain, C., & Potts, S. G. (2011) The potential impacts of insecticides on the life-history traits of bees and the consequences for pollination. *Basic and Applied Ecology*, 12(4), 321–331. doi:10.1016/j.baae.2010.12.004
- Bryden, J; Gill, RJ; Mitton, RAA; Raine, NE; Jansen, VAA (2013) Chronic sublethal stress causes bee colony failure. *ECOLOGY LETTERS* 16(12):1463-1469; 10.1111/ele.12188
- Cameron, S. a, Lozier, J. D., Strange, J. P., Koch, J. B., Cordes, N., Solter, L. F., & Griswold, T. L. (2011). Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences of the United States of America*, 108(2), 662–7. doi:10.1073/pnas.1014743108
- Cane J. H. and V. J. Tepedino (2001) Causes and extent of declines among native North American invertebrate pollinators: detection, evidence, and consequences. *Conservation Ecology* 5(1): 1 at <http://www.consecol.org/vol5/iss1/art1/>
- Castillo, C., Chen, H., Graves, C., Maisonnasse, A., Le Conte, Y., & Plettner, E. (2012) Biosynthesis of ethyl oleate, a primer pheromone, in the honey bee (*Apis mellifera* L.). *Insect biochemistry and molecular biology*, (March), 1–13. doi:10.1016/j.ibmb.2012.02.002
- Chauzat, M., & Faucon, J. (2007). Pesticide residues in beeswax samples collected from honey bee colonies (*Apis mellifera* L.) in France. *Pest Management Science*, 1106(December 2006), 1100–1106. doi:10.1002/ps
- Chauzat, M.-P., Carpentier, P., Martel, A.-C., Bougeard, S., Cougoule, N., Porta, P., Lachaize, J., et al. (2009) Influence of Pesticide Residues on Honey Bee (Hymenoptera: Apidae) Colony Health in France. *Environmental Entomology*, 38(3), 514–523. doi:10.1603/022.038.0302

- Ciarlo, T. J., Mullin, C. a., Frazier, J. L., & Schmehl, D. R. (2012). Learning Impairment in Honey Bees Caused by Agricultural Spray Adjuvants. (G. Smagghe, Ed.) PLoS ONE, 7(7), e40848. doi:10.1371/journal.pone.0040848
- Colin, M. E., Bonmatin, J. M., Moineau, I., Gaimon, C., Brun, S., & Vermandere, J. P. (2004) A method to quantify and analyze the foraging activity of honey bees: relevance to the sublethal effects induced by systemic insecticides. Archives of environmental contamination and toxicology, 47(3), 387–95. doi:10.1007/s00244-004-3052-y
- Colla, SR; Szabo, ND; Wagner, DL; Gall, LF; Kerr, JT (2013) Response to Stevens and Jenkins' pesticide impacts on bumblebees: a missing piece. CONSERVATION LETTERS 6(3):215-216; 10.1111/conl.12019
- Cox-Foster, D., Conlan, S., Holmes, E. C., Palacios, G., Evans, J. D., Moran, N. a, Quan, P.-L., et al. (2007) A metagenomic survey of microbes in honey bee colony collapse disorder. Science (New York, N.Y.), 318(5848), 283–7. doi:10.1126/science.1146498
- Cresswell, J. E. (2011) A meta-analysis of experiments testing the effects of a neonicotinoid insecticide (imidacloprid) on honey bees. Ecotoxicology (London, England), 20(1), 149–57. doi:10.1007/s10646-010-0566-0
- Cresswell, J. E., Desneux, N., & Vanengelsdorp, D. (2012) Dietary traces of neonicotinoid pesticides as a cause of population declines in honey bees: an evaluation by Hill's epidemiological criteria. Pest Management Science (January). doi:10.1002/ps.3290
- Cresswell, James E.; Page, Christopher J.; Uygun, Mehmet B.; Holmbergh, Marie; Li, Yueru; Wheeler, Jonathan G.; Laycock, Ian; Pook, Christopher J.; de Ibarra, Natalie Hempel; Smirnov, Nick; Tyler, Charles R. (2012) Differential sensitivity of honey bees and bumble bees to a dietary insecticide (imidacloprid) ZOOLOGY 115(6):365-371; 10.1016/j.zool.2012.05.003
- Cresswell, JE (2011) A meta-analysis of experiments testing the effects of a neonicotinoid insecticide (imidacloprid) on honey bees. Ecotoxicology 20(1):149-57. Epub 2010 Nov 16.
- Cutler, G. C., & Scott-dupree, C. D. (2007) Exposure to Clothianidin Seed-Treated Canola Has No Long-Term Impact on Honey Bees Exposure to Clothianidin Seed-Treated Canola Has No Long-Term Impact on Honey Bees. Journal of Economic Entomology, 100(3), 765–772. doi:10.1603/0022-0493(2007)100
- Dai P-L, Wang Q, Sun J-H, Liu F, Wang X, Wu Y-Y and Zhou T (2010) Effects of sub-lethal concentrations of bifenthrin and deltamethrin on fecundity, growth, and development of the honeybee *Apis mellifera ligustica*. Environmental Toxicology and Chemistry 29: 644–649.
- Decourtye and Devillers (2010) Ecotoxicity of Neonicotinoid Insecticides to Bees. In, Insect Nicotinic Acetylcholine Receptors, Advances in Experimental Medicine and Biology 683: 85-95, DOI: 10.1007/978-1-4419-6445-8_8.
- Decourtye, A (2003) Learning performances of honeybees (*Apis mellifera* L) are differentially affected by imidacloprid according to the season. Pest Manag Sci 59: 269-278.
- Decourtye, A. (2004) Imidacloprid impairs memory and brain metabolism in the honeybee (*Apis mellifera* L.). Pesticide Biochemistry and Physiology, 78(2), 83–92. doi:10.1016/j.pestbp.2003.10.001
- Decourtye, A., Devillers, J., Aupinel, P., Brun, F., Bagnis, C., Fourrier, J., & Gauthier, M. (2011) Honeybee tracking with microchips: a new methodology to measure the effects of pesticides. Ecotoxicology (London, England), 20(2), 429–37. doi:10.1007/s10646-011-0594-4
- Decourtye, A., Devillers, J., Cluzeau, S., Charreton, M., & Pham-Delègue, M.-H. (2004) Effects of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. Ecotoxicology and environmental safety, 57(3), 410–9. doi:10.1016/j.ecoenv.2003.08.001

- DeGrandi-Hoffman, G, et al (2010) The effect of diet on protein concentration, hypopharyngeal gland development and virus load in worker honey bees (*Apis mellifera* L.). *Journal of Insect Physiology* 56: 1184–1191.
- Derecka, K; Blythe, MJ; Malla, S; Genereux, DP; Guffanti, A; Pavan, P; Moles, A; Snart, C; Ryder, T; Ortori, CA; Barrett, DA; Schuster, E; Stoger, R(2013) Transient Exposure to Low Levels of Insecticide Affects Metabolic Networks of Honeybee Larvae. *PLOS ONE* 8(7):e68191; 10.1371/journal.pone.0068191
- Desneux, N., Decourtye, A., & Delpuech, J.-M. (2007) The sublethal effects of pesticides on beneficial arthropods. *Annual review of entomology*, 52, 81–106. doi:10.1146/annurev.ento.52.110405.091440
- Di Prisco, G; Cavaliere, V; Annoscia, D; Varricchio, P; Caprio, E; Nazzi, F; Gargiulo, G; Pennacchio, F (2013) Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA* 110(46): 18466-18471; 10.1073/pnas.1314923110
- Dittbrenner, N, et al (2011) Assessment of short and long-term effects of imidacloprid on the burrowing behaviour of two earthworm species (*Aporrectodea caliginosa* and *Lumbricus terrestris*) by using 2D and 3D post-exposure techniques. *Chemosphere* 84(10): 1349–1355.
- Dively, G. P., & A. Kamel (2012) Insecticide residues in pollen and nectar of a cucurbit crop and their potential exposure to pollinators. *Journal of Agricultural and Food Chemistry* 60:4449-4456.
- Donnarumma, L., Pulcini, P., Pochi, D., Rosati, S., Lusco, L., & Conte, E. (2011). Preliminary study on persistence in soil and residues in maize of imidacloprid. *Journal of environmental science and health. Part. B, Pesticides, food contaminants, and agricultural wastes*, 46(6), 469–72. doi:10.1080/03601234.2011.583848
- Easton, AH; Goulson, D (2013)The Neonicotinoid Insecticide Imidacloprid Repels Pollinating Flies and Beetles at Field-Realistic Concentrations. *PLOS ONE* 8(1) e54819; 10.1371/journal.pone.0054819
- Ebadi, R, et al (1980) Effects of carbon dioxide and low temperature narcosis on honey bees *Apis mellifera*. *Environmental Entomology* 9: 144–147.
- Eiri, D and JC Nieh (2012) A nicotinic acetylcholine receptor agonist affects honey bee sucrose responsiveness and decreases waggle dancing. *The Journal of Experimental Biology* 215: 2022-2029.
- El Hassani, A. K., Dacher, M., Gary, V., Lambin, M., Gauthier, M., & Armengaud, C. (2008) Effects of sublethal doses of acetamiprid and thiamethoxam on the behavior of the honeybee (*Apis mellifera*). *Archives of environmental contamination and toxicology*, 54(4), 653–61. doi:10.1007/s00244-007-9071-8
- Ellis, M. D. (2010) Pesticides Applied to Crops and Honey Bee Toxicity. *American Bee Journal*, 485-486. Retrieved from <http://entomology.unl.edu/faculty/ellispubs/CAP.pdf>
- Elston, C; Thompson, HM; Walters, KFA (2013) Sub-lethal effects of thiamethoxam, a neonicotinoid pesticide, and propiconazole, a DMI fungicide, on colony initiation in bumblebee (*Bombus terrestris*) micro-colonies. *APIDOLOGIE* 44(5):563-574; 10.1007/s13592-013-0206-9
- Faucon J.-P, Aurières C, Drajnudel P, Mathieu L, Ribière, M, Martel A-C, Zeggane S, et al. (2005) Experimental study on the toxicity of imidacloprid given in syrup to honey bee (*Apis mellifera*) colonies. *Pest management science*, 61(2), 111–25. doi:10.1002/ps.957
- Fernandez, F. C., Da Cruz-Landim, C., & Malaspina, O. (2012) Influence of the insecticide pyriproxyfen on the flight muscle differentiation of *Apis mellifera* (Hymenoptera, Apidae). *Microscopy research and technique*, 000(November 2011), 1–5. doi:10.1002/jemt.22003
- Frazier M, Mullin CA, Frazier J, & Ashcraft S (2008) What Have Pesticides Got to Do with It? *American Bee Journal*, 1–5.

Frost, E (2011) Effects of cold immobilization and recovery period on honeybee learning, memory, and responsiveness to sucrose. *Journal of Insect Physiology* 57: 1385–1390.

Gill, Richard J.; Ramos-Rodriguez, Oscar; Raine, Nigel E.(2012) Combined pesticide exposure severely affects individual- and colony-level traits in bees. *NATURE* 491(7422): 105-U119; 10.1038/nature11585

Girolami V, Mazzon L, Squartini A, Mori N, Marzaro M, Bernardo D, Greatti M, et al. (2009) Translocation of Neonicotinoid Insecticides from Coated Seeds to Seedling Guttation Drops: A Novel Way of Intoxication for Bees. *Journal of Economic Entomology*, 102(5), 1808–1815. doi:10.1603/029.102.0511

Girolami, V; Marzaro, M; Vivan, L; Mazzon, L; Giorio, C; Marton, D; Tapparo, A (2013) Aerial powdering of bees inside mobile cages and the extent of neonicotinoid cloud surrounding corn drillers. *JOURNAL OF APPLIED ENTOMOLOGY* 137(1-2):35-44; 10.1111/j.1439-0418.2012.01718.x

Giroud, B; Vauchez, A; Vulliet, E; Wiest, L; Bulete, A (2013) Trace level determination of pyrethroid and neonicotinoid insecticides in bee bread using acetonitrile-based extraction followed by analysis with ultra-high-performance liquid chromatography-tandem mass spectrometry. *JOURNAL OF CHROMATOGRAPHY A* 1316(53): 61; 10.1016/j.chroma.2013.09.088

Goulson, D (2013) REVIEW: An overview of the environmental risks posed by neonicotinoid insecticides. *JOURNAL OF APPLIED ECOLOGY* 50(4):977-987; 10.1111/1365-2664.12111

Gregorc A, Evans JD, Scharf M & Ellis JD (2012) Gene expression in honey bee (*Apis mellifera*) larvae exposed to pesticides and Varroa mites (*Varroa destructor*). *Journal of insect physiology*. Elsevier Ltd. doi:10.1016/j.jinsphys.2012.03.015

Guez D. (2003) Effects of imidacloprid metabolites on habituation in honeybees suggest the existence of two subtypes of nicotinic receptors differentially expressed during adult development. *Pharmacology Biochemistry and Behavior*, 75(1), 217–222. doi:10.1016/S0091-3057(03)00070-4

Haarmann T, Spivak M, Weaver D, Weaver B, & Glenn T. (2002) Effects of Fluvalinate and Coumaphos on Queen Honey Bees (Hymenoptera: Apidae) in Two Commercial Queen Rearing Operations. *Journal of Economic Entomology*, 95(1), 28–35. doi:10.1603/0022-0493-95.1.28

Hatjina, F; Papaefthimiou, C; Charistos, L; Dogaroglu, T; Bouga, M; Emmanouil, C; Arnold, G (2013) Sublethal doses of imidacloprid decreased size of hypopharyngeal glands and respiratory rhythm of honeybees in vivo. *APIDOLOGIE* 44(4):467-480; 10.1007/s13592-013-0199-4

Hawthorne DJ & Dively GP (2011). Killing Them with Kindness? In-Hive Medications May Inhibit Xenobiotic Efflux Transporters and Endanger Honey Bees. (G. Smagghe, Ed.) *PLoS ONE*, 6(11), e26796. doi:10.1371/journal.pone.0026796

Henry M, Reguin M., Requier F et al (2012) A common pesticide decreases foraging success and survival in honey bees. *Science* 336 (6079): 348-350.

Hoffmann E J, & SJ Castle (2012) Imidacloprid in melon guttation fluid: A potential mode of exposure for pest and beneficial organisms. *Journal of Economic Entomology* 105:67-71.

Hunt, Greg; Krupke, Christian (2012) Managed Pollinator CAP Coordinated Agricultural Project A National Research and Extension Initiative to Reverse Pollinator Decline Neonicotinoid Seed Treatments and Honey Bee Health *AMERICAN BEE JOURNAL* 152(9):889-891

Huntzinger ACI, James RR, Bosch J, & Kemp WP (2008) Fungicide Tests on Adult Alfalfa Leafcutting Bees (Hymenoptera: Megachilidae) *Journal of Economic Entomology*, 101(4), 1088–1094. doi:10.1603/0022-0493(2008)101

Iwasa, T. (2004) Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Protection*, 23(5), 371–378. doi:10.1016/j.cropro.2003.08.018

- James RR & Xu J (2012). Mechanisms by which pesticides affect insect immunity. *Journal of invertebrate pathology*, 109(2), 175–82. doi:10.1016/j.jip.2011.12.005
- Johnson, R. M., Ellis, M. D., Mullin, C. a., & Frazier. M. (2010) Pesticides and honey bee toxicity - USA. *Apidologie*. Retrieved from <http://entomology.unl.edu/faculty/ellispubs/Pesticides.pdf>
- Köhler, A., Pirk, C. W. W., & Nicolson, S. W. (2011) Honeybees and nectar nicotine: Deterrence and reduced survival versus potential health benefits. *Journal of insect physiology*, (December). doi:10.1016/j.jinsphys.2011.12.002
- Krischik, VA, Al Landmark, and GE. Heimpel (2007) Soil-Applied Imidacloprid Is Translocated to Nectar and Kills Nectar-Feeding *Anagyrus pseudococci* (Girault) (Hymenoptera: Encyrtidae). *Environ. Entomol.* 36(5): 1238-1245.
- Krupke, C. H., Hunt, G. J., Eitzer, B. D., Andino, G., & Given, K. (2012) Multiple Routes of Pesticide Exposure for Honey Bees Living Near Agricultural Fields. (G. Smagghe, Ed.) *PLOS ONE*, 7(1), e29268. doi:10.1371/journal.pone.0029268
- Lambin, M., Armengaud, C., Raymond, S., & Gauthier, M. (2001) Imidacloprid-induced facilitation of the proboscis extension reflex habituation in the honeybee. *Archives of insect biochemistry and physiology*, 48(3), 129–34. doi:10.1002/arch.1065
- Larson, JL; Redmond, CT; Potter, DA (2013) Assessing Insecticide Hazard to Bumble Bees Foraging on Flowering Weeds in Treated Lawns. *PLOS ONE* 8(6):e66375; 10.1371/journal.pone.0066375
- Laurino, D., Porporato, M., Patetta, A., & Manino, A. (2011) Toxicity of neonicotinoid insecticides to honey bees : laboratory tests. *Bulletin of Insectology*, 64(1), 107–113.
- Laurino, D; Manino, A; Patetta, A; Porporato, M (2013) Toxicity of neonicotinoid insecticides on different honey bee genotypes. *BULLETIN OF INSECTOLOGY* 66(1):119-126
- Laycock, I; Cresswell, JE(2013) Repression and Recuperation of Brood Production in *Bombus terrestris* Bumble Bees Exposed to a Pulse of the Neonicotinoid Pesticide Imidacloprid. *PLOS ONE* 8(11): e79872; 10.1371/journal.pone.0079872
- Laycock, Ian; Lenthall, Kate M.; Barratt, Andrew T.; Cresswell, James E.(2012) Effects of imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*). *ECOTOXICOLOGY* 21(7):1937-1945 ; 10.1007/s10646-012-0927-y
- Lu, C, KM Warchol, RA Callahan (2012) In situ replication of honey bee colony collapse disorder. *Bulletin of Insectology* 65 (1): 99-106.
- Lundin, Ola; Rundlof, Maj; Smith, Henrik G.; Bommarco, Riccardo (2012) Towards Integrated Pest Management in Red Clover Seed Production *JOURNAL OF ECONOMIC ENTOMOLOGY* 105(5):1620- 1628; 10.1603/EC12179
- Mao, W., Schuler, M. A., & Berenbaum, M. R. (2011) CYP9Q-mediated detoxification of acaricides in the honey bee (*Apis mellifera*). *Proceedings of the National Academy of Sciences*, 6–11. doi:10.1073/pnas.1109535108
- Martel, A.-C., Zeggane, S., Aurières, C., Drajnudel, P., Faucon, J.-P., & Aubert, M. (2007) Acaricide residues in honey and wax after treatment of honey bee colonies with Apivar or Asuntol 50. *Apidologie*, 38, 534–544. doi:10.1051/apido
- Marzaro, M., Vivan, L., Targa, A., Mazzon, L., Mori, N., Greatti, M., Toffolo, E. P., et al. (2011) Lethal aerial powdering of honey bees with neonicotinoids from fragments of maize seed coat. *Bulletin of Insectology*, 64(1), 119–126.
- Matsumoto, T(2013) Reduction in homing flights in the honey bee *Apis mellifera* after a sublethal dose of neonicotinoid insecticides. *BULLETIN OF INSECTOLOGY* 66 (1):1-9
- Maxim, L., & van der Sluijs, J. P. (2007). Uncertainty: Cause or effect of stakeholders' debates? Analysis of a case study: The risk for honeybees of the insecticide Gaucho((R)). *Science of the Total Environment*, 376, 1–17. doi:10.1016/j.scitotenv.2006.12.052

- Maxim, L., & van der Sluijs, J. P. (2009). Multicausality: from science to discourse, Evidence from a French case study. In A. P. Guimaraes & S. Funtowicz (Eds.), *Science for Policy* (pp. 67–80). Oxford: Oxford University Press. Retrieved from <http://www.ivds.nl/bijensterfte/03-Multi-Causal.pdf>
- Maxim, L., & van der Sluijs, J. P. (2010) Expert explanations of honeybee losses in areas of extensive agriculture in France: Gaucho (R) compared with other supposed causal factors. *Environmental Research Letters*, 5. doi:10.1088/1748-9326/5/1/014006
- Medrzycki P, Montanari R, Bortolotti L, Sabatini AG, Maini S, Porrini C. (2003) Effects of imidacloprid administered in sub-lethal doses on honey bee behaviour. Laboratory tests. *B Insectol* 56(1): 59–62.
- Medrzycki, P, et al (2010) Influence of brood rearing temperature on honey bee development and susceptibility to poisoning by pesticides. *Journal of Apicultural Research* 49(1): 52-59
- Mommaerts, V., Reynders, S., Boulet, J., Besard, L., Sterk, G., & Smagghe, G. (2010) Risk assessment for side-effects of neonicotinoids against bumblebees with and without impairing foraging behavior. *Ecotoxicology (London, England)*, 19(1), 207–15. doi:10.1007/s10646-009-0406-2
- Mullin, C. a., Frazier, M., Frazier, J. L., Ashcraft, S., Simonds, R., Vanengelsdorp, D., & Pettis, J. S. (2010) High levels of miticides and agrochemicals in North American apiaries: implications for honey bee health. *PloS one*, 5(3), e9754. doi:10.1371/journal.pone.0009754
- Mussen, Eric; Oliver, Randy (2012) Testing of Bee Feed Syrups for Neonicotinoid Residues *AMERICAN BEE JOURNAL* 152(8):803-804
- Nguyen, B. K., Saegerman, C., Pirard, C., Mignon, J., Widart, J., Thirionet, B., Verheggen, F. J., et al. (2009) Does Imidacloprid Seed-Treated Maize Have an Impact on Honey Bee Mortality? *Journal of Economic Entomology*, 102(2), 616–623. doi:10.1603/029.102.0220
- Nuyttens, D; Devarrewaere, W; Verboven, P; Foque, D (2013) Pesticide-laden dust emission and drift from treated seeds during seed drilling: a review. *PEST MANAGEMENT SCIENCE* 69(5):564-575; 10.1002/ps.3485
- Pareja, L., Colazzo, M., Pérez-Parada, A., Niell, S., Carrasco-Letelier, L., Besil, N., Cesio, M. V., et al. (2011) Detection of pesticides in active and depopulated beehives in Uruguay. *International journal of environmental research and public health*, 8(10), 3844–58. doi:10.3390/ijerph8103844
- Pérez-Parada, A., Colazzo, M., Besil, N., Geis-Asteggiate, L., Rey, F., & Heinzen, H. (2011) Determination of coumaphos, chlorpyrifos and ethion residues in propolis tinctures by matrix solid-phase dispersion and gas chromatography coupled to flame photometric and mass spectrometric detection. *Journal of chromatography. A*. doi:10.1016/j.chroma.2011.06.097
- Pettis, J. S., Collins, A., Wilbanks, R., & Feldlaufer, M. F. (2004) Effects of coumaphos on queen rearing in the honey bee, *Apis mellifera*. *Apidologie*, 35, 605–610. doi:10.1051/apido
- Pettis, JS, et al (2012) Pesticide exposure in honey bees results in increased levels of the gut pathogen Nosema. *Naturwissenschaften* 99(2):153-8. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3264871/?tool=pubmed>
- Pilling, E. D., & Jepson, P. C. (1993). Synergism between EBI fungicides and a pyrethroid insecticide in the honeybee (*Apis mellifera*). *Pesticide Science*, 39(4), 293–297. doi:10.1002/ps.2780390407
- Pilling, E; Campbell, P; Coulson, M; Ruddle, N; Tornier, I (2013) A Four-Year Field Program Investigating Long-Term Effects of Repeated Exposure of Honey Bee Colonies to Flowering Crops Treated with Thiamethoxam. *PLOS ONE* 8(10): e77193; 10.1371/journal.pone.0077193
- Ramakrishnan, R (1999) Imidacloprid-enhanced *Reticulitermes flavipes* (Kollar) (Isoptera: Rhinotermitidae) susceptibility to the entomopathogen *Metarhizium anisopliae* (Metsch.) Sorokin. *J. Econ. Entomol* 92:1125–1132.

- Ramirez-Romero, R., Chaufaux, J., & Pham-Delegue, M.-H. (2005) Effects of Cry1Ab protoxin , deltamethrin and imidacloprid on the foraging activity and the learning performances of the honeybee *Apis mellifera* , a comparative approach 1. *Apidologie*, 36(4), 601–611. doi:10.1051/apido
- Reetz, Jana E.; Zuehlke, Sebastian; Spittler, Michael; Wallner, Klaus(2012) A method for identifying water foraging bees by refractometer analysis: a spotlight on daily and seasonal water collecting activities of *Apis mellifera* L. *JOURNAL FÜR VERBRAUCHERSCHUTZ UND LEBENSMITTELSICHERHEIT-JOURNAL OF CONSUMER PROTECTION AND FOOD SAFETY* 7(4):283-290; 10.1007/s00003-012-0799-3
- Rortais, A., Aronld, G., Halm, M., & Touffet-Briens, F. (2005) Modes of honeybees exposure to systemic insecticides: estimated amounts of contaminated pollen and nectar consumed by different categories of bees. *Apidologie*, 36(1), 71–83. doi:10.1051/apido
- Rossi, CD; Roat, TC; Tavares, DA; Cintra-Socolowski, P; Malaspina, O (2013) Effects of sublethal doses of imidacloprid in malpighian tubules of africanized *Apis mellifera* (Hymenoptera, Apidae). *MICROSCOPY RESEARCH AND TECHNIQUE* 76(5):552-558; 10.1002/jemt.22199
- Sande, D., Mullen, J., Wetzstein, M., & Houston, J. (2011) Environmental impacts from pesticide use: a case study of soil fumigation in Florida tomato production. *International journal of environmental research and public health*, 8(12), 4649–61. doi:10.3390/ijerph8124649
- Schmuck, R (2004) Effects of a chronic dietary exposure of the honeybee *Apis mellifera* (Hymenoptera: Apidae) to imidacloprid. *Arch. Environ. Contam. Toxicol.* 47: 471–478.
- Schmuck, R., Schöning, R., Stork, A., & Schramel, O. (2001) Risk posed to honeybees (*Apis mellifera* L, Hymenoptera) by an imidacloprid seed dressing of sunflowers. *Pest management science*, 57(3), 225–38. doi:10.1002/ps.270
- Schneider CW, et al (2012) RFID tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of *Apis mellifera*. <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0030023>
- Scott-Dupree, C. D., Conroy, L., & Harris, C. R. (2009). Impact of Currently Used or Potentially Useful Insecticides for Canola Agroecosystems on *Bombus impatiens* (Hymenoptera: Apidae), *Megachile rotundata* (Hymenoptera: Megachilidae), and *Osmia lignaria* (Hymenoptera: Megachilidae). *Journal of Economic Entomology*, 102(1), 177–182. doi:10.1603/029.102.0125
- Sgolastra, Fabio; Renzi, Teresa; Draghetti, Stefano; Medrzycki, Piotr; Lodesani, Marco; Maini, Stefano; Porrini, Claudio (2012) Effects of neonicotinoid dust from maize seed-dressing on honey bees. *BULLETIN OF INSECTOLOGY* 65(2):273-280
- Smodis Skerl MI, Kmecl V, Gregorc A. (2010) Exposure to pesticides at sublethal level and their distribution within a honey bee (*Apis mellifera*) colony. *Bulletin of Environmental Contamination and Toxicology* 85(2): 125–8.
- Stadler T, et al (2003) Long-term toxicity assessment of imidacloprid to evaluate side effects on honey bees exposed to treated sunflower in Argentina, *Bull Insect* 2003; 56:77-81.
- Stoner KA and BD Eitzer (2012) movement of soil-applied imidacloprid and thiamethoxam into nectar and pollen of squash (*Cucurbita pepo*). *PLoS ONE* 7(6): e39114. doi:10.1371/journal.pone.0039114
- Suchail, S., De Sousa, G., Rahmani, R., & Belzunces, L. P. (2004) In vivo distribution and metabolism of ¹⁴C-imidacloprid in different compartments of *Apis mellifera* L. *Pest management science*, 60(11), 1056–62. doi:10.1002/ps.895
- Suchail, S., Debrauwer, L., & Belzunces, L. P. (2004) Metabolism of imidacloprid in *Apis mellifera*. *Pest management science*, 60(3), 291–6. doi:10.1002/ps.772
- Tang, Hongzhi; Li, Jian; Hu, Haiyang; Xu, Ping (2012) newly isolated strain of *Stenotrophomonas* sp hydrolyzes acetamiprid, a synthetic insecticide. *PROCESS BIOCHEMISTRY* 47(12):1820- 1825; 10.1016/j.procbio.2012.06.008

- Tapparo, A., Marton, D., Giorio, C., Zanella, A., Solda, L., Marzaro, M., Vivan, L., et al. (2012) Assessment of the Environmental Exposure of Honeybees to Particulate Matter Containing Neonicotinoid Insecticides Coming from Corn Coated Seeds. *Environmental Science and Technology*.
- Tapparo, A; Giorio, C; Solda, L; Bogialli, S; Marton, D; Marzaro, M; Girolami, V (2013) UHPLC-DAD method for the determination of neonicotinoid insecticides in single bees and its relevance in honeybee colony loss investigations. *ANALYTICAL AND BIOANALYTICAL CHEMISTRY* 405(2-3) 1007-1014; 10.1007/s00216-012-6338-3
- Tautz, J, et al (2003) Behavioral performance in adult honey bees is influenced by the temperature experienced during their pupal development. *Proc Natl Acad Sci* 100(12): 7343–7347.
- Tennekes, H (2010) The significance of the Druckrey–Küpfmüller equation for risk assessment—The toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology*. 2010 Sep 30;276(1):1-4. <http://www.ncbi.nlm.nih.gov/pubmed/20803795>
- Tennekes, H. (2011) The significance of the Druckrey–Küpfmüller equation for risk assessment—The toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time: Responding to a Letter to the Editor by Drs. C. Maus and R. Nauen of Bayer CropScience AG. *Toxicology Letters* 280(3), 173–175. doi:10.1016/j.tox.2010.11.015
- Tennekes, H. A., & Sánchez-bayo, F. (2011) Time-Dependent Toxicity of Neonicotinoids and Other Toxicants: Implications for a New Approach to Risk Assessment. *Environmental & Analytical Toxicology*, 1–8. doi:10.4172/2161-0525.S4-001
- Thompson, H. M., Wilkins, S., Battersby, A. H., Waite, R. J., & Wilkinson, D. (2007) Modelling long-term effects of IGRs (Insect Growth Regulators) on honey bee colonies. *Pest Management Science*, 63, 1081–1084. doi:10.1002/ps
- Tomé, H. V. V., Martins, G. F., Lima, M. A. P., Campos, L. A. O., & Guedes, R. N. C. (2012) Imidacloprid-Induced Impairment of Mushroom Bodies and Behavior of the Native Stingless Bee *Melipona quadrifasciata anthidioides*. *PLoS one*, 7(6), e38406. doi:10.1371/journal.pone.0038406
- Tomizawa, M and JE Casida (2009) Molecular recognition of neonicotinoid insecticides: the determinants of life or death. *Acc. Chem. Res.* 42(2): 260–269.
- Tsigouri, a. D., Menkissoglu-Spiroudi, U., Thrasyvoulou, A., & Diamantidis, G. (2004) Fluralinate Residues in Honey and Beeswax after Different Colony Treatments. *Bulletin of Environmental Contamination and Toxicology*, 72(5), 975–982. doi:10.1007/s00128-004-0339-7
- Tsiropoulos, N. G., & Amvrazi, E. G. (2011) Determination of pesticide residues in honey by single-drop microextraction and gas chromatography. *Journal of AOAC International*, 94(2), 634–44. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21563700>
- Underwood, R. M., & Currie, R. W. (2003) The effects of temperature and dose of formic acid on treatment efficacy against *Varroa destructor* (Acari: Varroidae), a parasite of *Apis mellifera* (Hymenoptera: Apidae). *Experimental & applied acarology*, 29(3-4), 303–13. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14635816>
- Van Dijk, TC; Van Staalduinen, MA; Van der Sluijs, JP. (2013) Macro-Invertebrate Decline in Surface Water Polluted with Imidacloprid. *PLOS ONE* 8(5): e62374; 10.1371/journal.pone.0062374
- vanEngelsdorp D, Evans JD, Donovall L, Mullin C, Frazier M, Frazier J, et al. (2009) “Entombed pollen”: a new condition in honey bee colonies associated with increased risk of colony mortality. *J Invertebr Pathol* 101(2): 147–149.
- vanEngelsdorp, D., Evans, J. D., Saegerman, C., Mullin, C. a., Haubruge, E., Nguyen, B. K., Frazier, M., et al. (2009) Colony collapse disorder: a descriptive study. *PLoS one*, 4(8), e6481. doi:10.1371/journal.pone.0006481

vanEngelsdorp, D., Speybroeck, N., Evans, J. D., Nguyen, B. K., Mullin, C. a., Frazier, M., Frazier, J., et al. (2010) Weighing Risk Factors Associated with Bee Colony Collapse Disorder by Classification and Regression Tree Analysis. *Journal of Economic Entomology*, 103(5), 1517–1523. doi:10.1603/EC09429

Vidau C, Diogon M, Aufauvre J, Fontbonne R, Viguès B, et al. (2011) Exposure to sublethal doses of fipronil and thiacloprid highly increases mortality of honey bees previously infected by *Nosema ceranae*. *PLoS ONE* 6(6): e21550.

Vidau, C., Diogon, M., Aufauvre, J., Fontbonne, R., Viguès, B., Brunet, J.-L., Texier, C., et al. (2011) Exposure to Sublethal Doses of Fipronil and Thiacloprid Highly Increases Mortality of Honeybees Previously Infected by *Nosema ceranae*. (E. Didier, Ed.) *PLoS ONE*, 6(6), 8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21738706>

Wang, R., Wang, Z., Yang, H., Wang, Y., & Deng, A. (2011). Highly sensitive and specific detection of neonicotinoid insecticide imidacloprid in environmental and food samples by a polyclonal antibody-based enzyme-linked immunosorbent assay. *Journal of the Science of Food and Agriculture*, (September). doi:10.1002/jsfa.4691

Whitehorn, P, et al (2012) Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* 336 (6079): 351-352.

Williamson, SM; Baker, DD; Wright, GA (2013) Acute exposure to a sublethal dose of imidacloprid and coumaphos enhances olfactory learning and memory in the honeybee *Apis mellifera* *INVERTEBRATE NEUROSCIENCE* 63(70):10.1007/s10158-012-0144-7

Wu, J. Y., Anelli, C. M., & Sheppard, W. S. (2011) Sub-Lethal Effects of Pesticide Residues in Brood Comb on Worker Honey Bee (*Apis mellifera*) Development and Longevity. (F. Marion-Poll, Ed.) *PLoS ONE*, 6(2), e14720. doi:10.1371/journal.pone.0014720

Wu, J. Y., Smart, M. D., Anelli, C. M., & Sheppard, W. S. (2012) Honey bees (*Apis mellifera*) reared in brood combs containing high levels of pesticide residues exhibit increased susceptibility to *Nosema* (Microsporidia) infection. *Journal of invertebrate pathology*, 109(3), 326–9. doi:10.1016/j.jip.2012.01.005

Yanez, KP; Bernal, JL; Nozal, MJ; Martin, MT; Bernal, J (2013) Determination of seven neonicotinoid insecticides in beeswax by liquid chromatography coupled to electrospray-mass spectrometry using a fused-core column. *JOURNAL OF CHROMATOGRAPHY A* 1285:110-117; .1016/j.chroma.2013.02.032

Yang EC, Chuang YC, Chen YL, Chang LH. (2008) Abnormal foraging behavior induced by sublethal dosage of imidacloprid in the honey bee (Hymenoptera: Apidae). *J of Econ Entomol* 101(6): 1743-1748. doi:10.1603/0022-0493-101.6.1743

Zhang, J., Gao, H., Peng, B., Li, S., & Zhou, Z. (2011) Comparison of the performance of conventional, temperature-controlled, and ultrasound-assisted ionic liquid dispersive liquid-liquid microextraction combined with high-performance liquid chromatography in analyzing pyrethroid pesticides in honey samples. *Journal of Chromatography. A*. doi:10.1016/j.chroma.2011.07.102

Proceedings from Symposia

Alix A, Chauzat MP, Duchard S, Lewis, G, Maus C, Miles MJ, Pilling E, Thompson HM, and Wallner K. (2009) Guidance for the assessment of risks to bees from the use of plant protection products applied as seed coating and soil applications – conclusions of the ICPBR dedicated working group. In: Oomen, PA and HM Thompson (Editors) (2009) *Hazards of Pesticides to Bees: International Commission for Plant-Bee Relationships, Bee Protection Group, 10th International Symposium*. http://www.jki.bund.de/fileadmin/dam_uploads/_veroeff/JKI_Archiv/JKI_Archiv_423.pdf.

Bonmatin, J.M., Marchand P.A, Charvet R, Colin M.E, (1994)“Fate of systemic insecticides in fields (Imidacloprid and Fipronil) and risks for pollinators, in First European Conference of Apidology”. Udine 19-23 September 2004.

Dively, GP, et al (2010) Sublethal and synergistic effects of pesticides
<http://agresearch.umd.edu/recs/WREC/files/2010Programs/EASSubletha%20Effects2010.pdf>

Hatjina, F and T Dogaroglu (2010) Imidacloprid effect on honey bees under laboratory conditions using hoarding cages. http://www.coloss.org/publications/proceedings_workshop_bologna_2010

Pimentel, D. (2001) Environmental effects of pesticides on public health, birds and other organisms. Rachel Carson and the Conservation Movement: Past Present and Future. Conference presented 10–12 August 2001, Shepherdstown, W.V. <http://rachels-carson-of-today.blogspot.com/2011/02/environmental-effects-of-pesticides-on.html>

Government, University, Industry, and NGO documents

AFSSA (2009) Mortalités, effondrements et affaiblissements des colonies d'abeilles (Weakening, collapse and mortality of bee colonies). <http://www.afssa.fr/Documents/SANT-Ra-MortaliteAbeilles.pdf>.

ApeNet. (2011) Effects of coated maize seed on honey bees - Report based on results obtained from the second year (2010) activity of the APENET project.

Are bees addicted to caffeine and nicotine? ScienceDaily. Retrieved February 8, 2011, from <http://www.sciencedaily.com/releases/2010/02/100210101504.htm>

California Department of Pesticide Regulation. Summary of Pesticide Use Report Data 2010 Indexed by Commodity <http://www.cdpr.ca.gov/docs/pur/pur10rep/comrpt10.pdf>

DEFRA (2007) Assessment of the Risk Posed to Honeybees by Systemic Pesticides, PS2322. Central Science Laboratory, (GB).
<http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=13502>

EFSA (2012) Statement on the findings in recent studies investigating sub-lethal effects in bees of some neonicotinoids in consideration of the uses currently authorised in Europe.
<http://www.efsa.europa.eu/fr/efsajournal/doc/2752.pdf>

EPA (2003) Data evaluation record honey bee - Acute oral LD50 test
http://www.epa.gov/pesticides/chem_search/cleared_reviews/csr_PC-044309_20-Mar-03_d.pdf

EPA (2005) EFED Registration Chapter for Clothianidin for use on Potatoes and Grapes as a spray treatment and as a Seed Treatment for Sorghum and Cotton.
http://www.epa.gov/pesticides/chem_search/cleared_reviews/csr_PC-044309_28-Sep-05_a.pdf

EPPO (2003) EPPO Standards: Environmental risk assessment scheme for plant protection products, Chapter 10, Honeybees. OEPP/EPPO Bulletin 33: 99–101.
[http://archives.eppo.int/EPPOStandards/PP3_ERA/pp3-10\(2\).pdf?utm_source=archives.eppo.org&utm_medium=int_redirect](http://archives.eppo.int/EPPOStandards/PP3_ERA/pp3-10(2).pdf?utm_source=archives.eppo.org&utm_medium=int_redirect)

EPPO (2010) Environmental risk assessment scheme for plant protection products. Chapter 10: honeybees. <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2338.2010.02419.x/pdf>

Hatjina, F and T Dogaroglu (2010) Imidacloprid effect on honey bees under laboratory conditions using hoarding cages. http://www.coloss.org/publications/proceedings_workshop_bologna_2010

Hopwood J, Mace Vaughan M, Shepherd M, Biddinger D, Mader E, Hoffman Black S, Mazzacano C. (2012) Are Neonicotinoids Killing Bees? (http://www.xerces.org/wp-content/uploads/2012/03/Are-Neonicotinoids-Killing-Bees_Xerces-Society1.pdf)

Lodesani, M, et al (2009) Effects of coated maize seed on honey bees: Effects on the brood.
http://www.cra-api.it/online/immagini/Apenet_2009_eng.pdf

Munkvold, G (2006) Seed Treatment <http://www.extension.iastate.edu/Publications/CS16.pdf>

Pesticide Risk Assessment for Pollinators: Summary of a SETAC Pellston Workshop
http://www.setac.org/sites/default/files/executivesummarypollinators_20sep2011.pdf

Response to petition <http://www.regulations.gov/#!documentDetail;D=EPA-HQ-OPP-2012-0334-0006> Technical supporting documents <http://www.regulations.gov/#!documentDetail;D=EPA-HQ-OPP-2012-0334-0012>

The Beekeepers Quarterly June 2012 (U.K.) Neonicotinoids—Our toxic countryside http://www.boerenlandvogels.nl/sites/default/files/BKQ%20108%20-%20Neonicotinoid%20Pesticides_0.pdf