

SELECTING

PLANTS

FOR

POLLINATORS



A GUIDE FOR GARDENERS, FARMERS, AND LAND MANAGERS IN THE



# ST. LAWRENCE LOWLANDS ECOREGION

QUEBEC CITY,

MONTREAL, TROIS-RIVIÈRES,

SAINT-HYACINTHE,

CORNWALL, BROCKVILLE,

OTTAWA, GATINEAU,

AND PEMBROKE



## TABLE OF CONTENTS

WHY SUPPORT POLLINATORS?	4
GETTING STARTED	5
ST. LAWRENCE LOWLANDS	(
MEET THE POLLINATORS	8
PLANT TRAITS	10
DEVELOPING PLANTINGS	12
FARMS	13
PUBLIC LANDS	14
HOME LANDSCAPES	15
PLANTS THAT ATTRACT POLLINATORS	16
HABITAT HINTS	20
HABITAT AND NESTING REQUIREMENTS	2
S.H.A.R.E.	22
CHECKLIST	22
RESOURCES AND FEEDBACK	23

This is one of several guides for different regions of North America. We welcome your feedback to assist us in making the future guides useful. Please contact us at

feedback@pollinator.org

### **SELECTING PLANTS FOR POLLINATORS**

A GUIDE FOR
GARDENERS, FARMERS,
AND LAND MANAGERS

IN THE

ST. LAWRENCE LOWLANDS

ECOREGION



QUEBEC CITY, MONTREAL,

TROIS-RIVIÈRES, SAINT-HYACINTHE,

CORNWALL, BROCKVILLE,

OTTAWA, GATINEAU,

AND PEMBROKE

A NAPPC AND POLLINATOR PARTNERSHIP CANADAM PLIBLICATION

Funding for this ecoregional guide was provided by The Stanley Smith Horticultural Trust

### WHY SUPPORT POLLINATORS?

IN THEIR 1996 BOOK, *The Forgotten Pollinators*, Buchmann and Nabhan estimated that animal pollinators are needed for the reproduction of 90% of flowering plants and one third of human food crops. Each of us depends on these industrious pollinators in a practical way to provide us with the wide range of foods we eat. In addition, pollinators are part of the intricate web that supports the biological diversity in natural ecosystems that helps sustain our quality of life.

Abundant and healthy populations of pollinators can improve fruit set and quality, and increase fruit size. In farming situations this increases production per hectare. In the wild, biodiversity increases and wildlife food sources increase. Apples, melons, raspberries, strawberries, peas, pumpkins, and tomatoes are some of the crop types grown in the St. Lawrence Lowlands ecoregion that rely on honey bees and native bees for pollination. Honey bees pollinate more than \$1 billion worth of crops in Canada each year.

Unfortunately, the numbers of many native pollinators are declining. They are threatened by habitat loss, disease, climate change, and the excessive and inappropriate use of pesticides. As a result research and conservation actions supporting native pollinators are increasingly active. Honey bee colony losses have significantly impacted beekeepers. Parasites, disease, pesticide use, insufficient nutrition, and transportation practices all impact honey bee health, and this in turn can impact the commercial pollination services honey bees provide. The efforts to understand the threats to commercial bees should help us understand other pollinators and their roles in the environment as well.

It is imperative that we take immediate steps to help pollinator populations thrive. The beauty of the situation is that by supporting pollinators' need for habitat, we support our own needs for food and support diversity in the natural world.

Thank you for taking time to consult this guide. By adding plants to your landscape that provide food and shelter for pollinators throughout their active seasons and by adopting pollinator friendly landscape practices, you can make a difference to both the pollinators and the people that rely on them.

Val Del

Val Dolcini President & CEO Pollinator Partnership

FLOWERING PLANTS **ACROSS WILD.** FARMED AND EVEN URBAN LANDSCAPES **ACTUALLY FEED THE** TERRESTRIAL WORLD. AND POLLINATORS **ARE THE GREAT CONNECTORS WHO ENABLE THIS GIANT FOOD SYSTEM TO** WORK FOR ALL WHO

ROGER LANG, CHAIRMAN,
 POLLINATOR PARTNERSHIP

EAT... INCLUDING US.







THIS REGIONAL GUIDE IS JUST ONE in a series of plant selection tools designed to provide you with information on how to plant local native plants for pollinators. Each of us can have a positive impact by providing the essential habitat requirements for pollinators including food, water, shelter, and enough space to raise their young.

Pollinators travel through the landscape without regard to property ownership or provincial boundaries. We've chosen to use the ecoregional classification system and to underscore the connections between climate and vegetation types that affect the diversity of pollinators in the environment.

The Canadian ecoregions are based on the National Ecological

Framework Report. The National Ecological Framework for Canada is a system created as a management tool and is used to predict responses to land management practices throughout large areas. This guide addresses pollinator-friendly land management practices in what is known as the St. Lawrence Lowlands ecoregion.

The seasonal cycle of sun and shade within the forests has created a changing pattern of bloom time for food plants and shelter needs for foraging, nesting, and migrating pollinators. Farms and residential areas provide a diverse range of soil types in both sunny and shady areas. With this diversity of locations many different species of plants may be used to improve pollinator habitats where they are lacking.

Long before there were homes and farms in the St. Lawrence Lowlands ecoregion, natural vegetation provided essential opportunities for wildlife, including pollinators. Farmers, land managers, and gardeners in this region have a wide palette of plants to use in the landscape. In choosing plants, aim to create habitat for pollinators that allow adequate food, shelter, and water sources. Most pollinators have very small home ranges. You will make a difference by understanding the vegetation patterns on the farm, forest, or neighbour's yard adjacent to your property. With this information in hand, your planting choices will better support the pollinators' need for food and shelter as they move through the landscape.

# UNDERSTANDING THE ST. LAWRENCE LOWLANDS



- The St. Lawrence Lowlands ecoregion makes up the eastern portion of the Mixed Wood Plains ecozone, and includes the following ecoregions: Manitoulin-Lake Simcoe, Frontenac Axis, and the Lake Erie Lowlands. The geographic location, watershed, and combination of gentle topography, fertile soils, warm growing seasons and abundant rainfall have made this area intensely farmed and populated.
- Not sure about which region you live or work in? Go to www. pollinator.org and click on Ecoregion Locator for help.
- 🔀 46,000 square kilometres in Québec.
- The topography includes till plains, lake plain, and rock landscapes.
- **%** Elevations ranging from 15 to 150 meters above sea level.
- ₩ The mean summer temperature is 16.5°C and the mean winter temperature is -7°C.
- The mean annual precipitation ranges 800-1000 mm.
- Favourable climate patterns in the region support a strong agricultural economy. More than 60% of the ecoregion is classified as dependable agricultural land.

#### **CHARACTERISTICS**

- **%** Dominant land cover is agriculture with significant areas of mixed forest and large urban centers.
- Mone of the highest areas of biodiversity found in Canada.
- **%** Dissected by population centres, farms, and roads.





#### St. Lawrence Lowlands includes:

Quebec City,

Montreal.

Trois-Rivières,

Saint-Hyacinthe,

Cornwall,

Brockville,

Ottawa, Gatineau,

and Pembroke



### MEET THE POLLINATORS



Ruby-throated Hummingbird, a summer species in the St. Lawrence Lowlands ecoregion.

#### Eastern Tiger Swallowtail.



# WHO ARE THE POLLINATORS?

#### BEES

Bees are the best documented pollinators in the natural and agricultural landscapes of the St. Lawrence Lowlands ecoregion. A wide range of plants in the Aster and Rose Families, apples, strawberries and raspberries are just a few plants that benefit from bee pollinators.

Most of us are familiar with the colonies of honey bees that have been the workhorses of agricultural pollination for years in Canada. They were imported from Europe almost 400 years ago and continue to be managed for honey production and pollination services. There are over 800 species of native ground and twig nesting bees in Canada. Most of these bee species live a solitary life; a minority are social and form colonies or nest in aggregations.

Native bees visit and pollinate many crops; in many cases they are better at transferring pollen than honey bees. Our native bees can be encouraged to do more to support agricultural endeavours if their needs for nesting habitat are met and if suitable sources of nectar, pollen, and water are provided. Bees come in a variety of body shapes and sizes, and even have tongues of different lengths. Native bees visit the widest range of flowers and crops of any pollinator group.

Bumble bees (Bombus spp.) form small colonies, usually underground making use of old rodent burrows or dense thatches. They are generalists, feeding on a wide range of plant types from May to September and are important pollinators of tomatoes and blueberries.

Sweat bees (family Halictidae) are medium to small-sized, slender bees that commonly nest underground. Various species are solitary while others form loose colonies, nesting side-by-side. Other common solitary bees include carpenter bees (Xylocopa virginica), which nest by chewing into wood; alkali bees (Andrena spp.), which nest underground and are common in the spring; leaf-cutter bees (Megachile spp.), which prefer dead trees or branches for their nest sites; and mason bees (Osmia spp.), which utilize cavities they find in stems and dead wood that they fill with mud.

#### **BUTTERFLIES**

Butterflies prefer open and sunny areas such as meadows and along woodland edges that provide bright flowers, water sources, and specific host plants for their caterpillars. Gardeners have been attracting butterflies to their gardens for some time. To encourage butterflies place flowering plants where they have full sun and are protected from the wind. They usually look for flowers that provide a good landing platform. Butterflies need open areas (e.g., bare earth, large stones) where they can bask, and moist soil from which they wick needed minerals. Butterflies eat rotten fruit and even dung, so don't clean up all the messes in your garden! By providing a safe place to eat and nest, gardeners can also support the pollination role that butterflies play in the landscape. In the Manitoulin-Lake Simcoe ecoregion it is common to see the Black Swallowtail (*Papilio polyxenes*), the Clouded Sulphur (Colias philodice), the Coral Hairstreak (Satyrium titus) and the Red Admiral (Vanessa atalanta).



#### **MOTHS**

Moths are most easily distinguished from butterflies by their antennae. Butterfly antennae are simple with a swelling at the end. Moth antennae differ from simple to featherlike, but never have a swelling at the tip. Butterfly bodies are not very hairy, while moth bodies are quite hairy and much more stout. In addition, butterflies typically are active during the day; moths at night. They are attracted to flowers that are strongly sweet smelling, open in late afternoon or night, and are typically white or pale coloured.

#### **BEETLES**

There are more than 9000 species of beetles in Canada and many of them can be found easily by looking inside flower heads. Gardeners have yet to intentionally draw beetles to their gardens, possibly because beetle watching isn't as inspiring as butterfly or bird watching. Yet beetles do play a role in pollination. Some have a bad reputation because they can leave a mess behind, damaging plant parts as they eat pollen. Beetle pollinated plants tend to be large, strong scented flowers and have the anthers and stigma exposed. Beetles are known to pollinate magnolia, paw paws, and yellow pond lilies.

#### **FLIES**

It may be hard to imagine why one would want to attract flies to the garden. However, flies are one of the most diverse group of pollinators. They include colouful flower flies and hover flies (*Syrphidae*), active bee flies (*Bombyliidae*), and tiny midges

that visit many plant species. Like bees, flies are hairy and can easily transport pollen from flower to flower. Flies primarily pollinate small flowers that bloom under shade and in seasonally moist habitats, but are also economically important as pollinators for a range of annual and bulbous ornamental flowers. Plants pollinated by flies include the American pawpaw (Asimina triloba), skunk cabbage (Symplocarpus foetiðus), goldenrod (Solidago spp.), and members of the carrot.

#### **BIRDS**

Hummingbirds are the primary birds that play a role in pollination in North America. Their long beaks and tongues draw nectar from tubular flowers. Pollen is carried on both their beaks and feathers. Regions closer to the tropics, with warmer climates, boast the largest number of hummingbird species and the greatest number of native plants to support their need for food. The Ruby-throated Hummingbird (*Archilochus colubris*) is the only common species in Quebec.

Hummingbirds can see the colour red; bees cannot. Many tropical flowers, grown as annuals in the Lake Erie Lowlands, along with native woodland edge plants, attract hummingbirds.

#### **BATS**

Though bat species in Canada are not pollinators, bats in the south western United States and Mexico are important pollinators of agave and cactus. The head shape and long tongues of nectar bats allows them to delve into flower blossoms and extract both pollen and nectar; pollen covers their hairy bodies and is transfer as they move from plant to plant.

#### **SPECIES AT RISK**

Species at Risk include endangered species, such as the Rusty-patched bumble bee, as well as threatened and special concern species. Other species are of conservation concern because of their rarity in Ontario and Quebec, but their formal conservation status has yet to be determined. Because habitat loss is one of the largest threats facing pollinators, the very best thing you can do is plant for them.

#### Rusty-patched bumble bee

- Though once widespread in Quebec, the only Canadian sightings of this bee since 2002 have been at The Pinery Provincial Park on Lake Huron.
- Bumble bees perform a special type of pollination called buzz pollination. They grab onto the flower and rapidly vibrate their flight muscles, dislodging pollen. Some plants, such as tomatoes, peppers, and cranberries greatly benefit from buzz pollination.
- While the exact cause of their steep decline are unknown, threats to Rusty-patched bumble bee populations include: pesticide use, habitat loss, and disease.

### PLANT TRAITS



# WHICH FLOWERS DO THE POLLINATORS PREFER?

NOT ALL POLLINATORS ARE found in each province, and some are more important in different parts of Canada. Use this page as a resource to understand the plants and pollinators where you live.

Plants can be grouped together based on the similar characteristics of their flowers. These floral characteristics can be useful to predict the type of pollination method or animal that is most effective for that group of plants. This association between floral characteristics and pollination method is called a pollination syndrome.

The interactions of animal pollinators and plants have influenced the evolution of both groups of organisms. A mutualistic relationship between the pollinator and the plant species helps the pollinator find necessary pollen and nectar sources and helps the plant reproduce by ensuring that pollen is carried from one flower to another.

Plant Trait	Pollinator								
	Bees	Beetles	Birds						
Colour	Bright white, yellow, blue, or UV	Dull white or green	Scarlet, orange, red or white						
Nectar Guides	Present	Absent	Absent						
Odour	Fresh, mild, pleasant	None to strongly fruity or fetid	None						
Nectar	Usually present	Sometimes present; not hidden	Ample; deeply hidden						
Pollen	Limited; often sticky and scented	Ample	Modest						
Flower Shape	Shallow; have landing platform; tubular	Large bowl-like, Magnolia	Large funnel like; cups, strong perch support						

This chart and more information on pollinator syndromes can be found at:



# AND THE POLLINATORS THEY ATTRACT

# **Pollinator**

Butterflies	Flies	Moths	Wind	
Bright, including red and purple	Pale and dull to dark brown or purple; flecked with translucent patches	Pale and dull red, purple, pink or white	Dull green, brown, or colourless; petals absent or reduced	
Present	Absent	Absent	Absent	
Faint but fresh	Putrid	Strong sweet; emitted at night	None	
Ample; deeply hidden	Usually absent	Ample; deeply hidden	None	
Limited	Modest in amount	Limited	Abundant; small, smooth, and not sticky	
Narrow tube with spur; wide landing pad	Shallow; funnel like or complex and trap-like	Regular; tubular without a lip	Regular; small and stigmas exerted	

http://www.fs.fed.us/wildflowers/pollinators/syndromes.shtml

# DEVELOPING LANDSCAPE PLANTINGS THAT PROVIDE POLLINATOR HABITAT

WHETHER YOU ARE A FARMER of many hectares, land manager of a large tract of land, or a gardener with a small lot, you can increase the number of pollinators in your area by making conscious choices to include plants that provide essential habitat for bees, butterflies, moths, beetles, hummingbirds and other pollinators.

#### FOOD:

Flowers provide nectar (high in sugar and necessary amino acids) and pollen (high in protein) to pollinators.

Fermenting fallen fruits also provide food for bees, beetles and butterflies. Specific plants, known as host plants, are eaten by the larvae of pollinators such as butterflies.

- Plant in groups to increase pollination efficiency. If a pollinator can visit the same type of flower over and over, it doesn't have to relearn how to enter the flower and can transfer pollen to the same species, instead of squandering the pollen on unreceptive flowers.
- Plant with bloom season in mind, providing food from early spring to late fall. (see Bloom Periods pp.16-19)
- Plant a diversity of plants to support a variety of pollinators. Flowers of different colour, fragrance, and season of bloom on plants of different heights will attract different pollinator species and provide pollen and nectar throughout the seasons.
- Many herbs and annuals, although not native, are very good for pollinators. Mint, oregano, garlic,

chives, parsley and lavender are just a few herbs that can be planted. Old fashioned zinnias, cosmos, and single sunflowers support bees and butterflies. Non-native crops, herbs, and ornamental flowers should only be planted in gardens.

- Recognize weeds that might be a good source of food. For example, dandelions provide nectar in the early spring before other flowers open. Milkweed is a host for the Monarch butterfly.
- Learn and utilize Integrated Pest Management (IPM) practices to address pest concerns. Minimize or eliminate the use of pesticides.

#### SHELTER:

Pollinators need protection from severe weather and from predators as well as sites for nesting and roosting.

- Incorporate different canopy layers in the landscape by planting trees, shrubs, and different-sized perennial plants.
- Leave dead snags for nesting sites of bees, and other dead plants and leaf litter for shelter.
- Avoid applying thick layers of mulch that are hard to dig through.
- Build bee boxes to encourage solitary, non-aggressive bees to nest on your property.
- Ground nesting bees are also attracted to lawns and short grass areas, especially if there is a south facing slope.
- Leave some areas of soil uncovered to provide ground nesting insects easy access to underground tunnels.
- Group plantings so that pollinators

can move safely through the landscape protected from predators.

• Include plants that are needed by butterflies during their larval development.

### **WATER:**

A clean, reliable source of water is essential to pollinators.

- Natural and human-made water features such as running water, pools, ponds, and small containers of water provide drinking and bathing opportunities for pollinators.
- Ensure the water sources have a shallow or sloping side so the pollinators can easily approach the water without drowning. Your current landscape probably includes many of these elements. Observe wildlife activity in your farm fields, woodlands, and gardens to determine what actions you can take to encourage other pollinators to feed and nest. Evaluate the placement of individual plants and water sources and use your knowledge of specific pollinator needs to guide your choice and placement of additional plants and other habitat elements. Minor changes by many individuals can positively impact the pollinator populations in your area. Watch for - and enjoy - the changes in your landscape!
- CAUTION LAND MANAGERS: Remember that pesticides are largely toxic to pollinators. Extreme caution is warranted if you choose to use any pesticide. Strategically apply pesticides only for problematic target species.



Apples, melons, raspberries, strawberries, peas, pumpkins, and tomatoes are a few of the food crops in the St. Lawrence Lowlands that will benefit from strong native bee populations that boost pollination efficiency. Incorporate different plants throughout the farm that provide food for native populations when targeted crops are not in flower.

Farmers have many opportunities to incorporate pollinator-friendly land management practices on their land which will benefit the farmer in achieving his or her production goals:

• Minimize the use of pesticides to reduce the impact on native pollinators. Spray when bees aren't active (just before dawn) and choose targeted ingredients.

- Be conscientious of the fact that different bees forage at different times of day. *Peponapsis pruinosa*, the squash bee, is active from dawn until noon. In the case of squash production, the best time to spray is in the evening rather than the early morning.
- Carefully consider the use of herbicides. Some weeds can provide needed food for pollinators.
- Minimize tillage to protect ground nesting pollinators.
- Consider creating designated permanently untilled areas for ground nesting bees along internal farm laneways.
- Ensure water sources are scattered throughout the landscape.
- Choose a variety of native plants to act as windbreaks, riparian buffers,

- and field borders throughout the farm.
- Plant unused areas of the farm with temporary cover crops that can provide food or with a variety of trees, shrubs, and flowers that provide both food and shelter for pollinators.
- Check with local not-for-profit organizations, local Conservation Authority or Ministry of Natural Resources and Forestry offices to see what technical support might be available to assist you in your effort to provide nectar, pollen, and larval food sources for pollinators on your farm.



### **PUBLIC LANDS**





Public lands are maintained for specific reasons ranging from high impact recreation to conservation. In the Manitoulin-Lake Simcoe, forests have been cut to allow for roads, energy infrastructure, buildings, open lawn areas, boat ramps, and vistas. These other lands can provide benefits to pollinator when managed correctly. Pollinators are encouraged by timed mowing, opening tree canopies, and planting of native flowers. Less disturbed natural areas can be augmented with plantings of native plant species. Existing plantings around buildings and parking areas should be evaluated to determine if pollinatorfriendly plants can be substituted or added to attract and support pollinators. Public land managers have a unique opportunity to use their plantings as an education tool to help others understand

the importance of pollinators in the environment through signs, brochures, and public programs. In an effort to increase populations of pollinators the land manager can:

- Inventory and become knowledgeable of local pollinators.
- Provide connectivity between vegetation areas by creating corridors of perennials, shrubs, and trees that provide pollinators shelter and food as they move through the landscape.
- Maintain a minimum of lawn areas that support recreational needs.
- Restrict the use of insecticides and herbicides.
- Provide water sources in large open areas.
- Maintain natural meadows and openings that provide habitats for sun-loving wildflowers and grasses.
- Remove invasive species.





Gardeners have a wide array of plants to use in their gardens. Native plants, plants introduced from years of plant exploration from around the world, and plants developed by professional and amateur breeders can be found in garden centres, in catalogs, and on web-sites. Use your knowledge of pollinator needs to guide your choices.

- Choose a variety of plants that will provide nectar and pollen throughout the growing season.
- Resist the urge to have a totally manicured lawn and garden. Leave bare ground for ground nesting bees. Leave areas of dead wood and leaf litter for other insects.
- Find local resources to help you in your efforts. Contact your local garden club or local Conservation Authority. Visit your regional botanic gardens and arboreta.

The scale of your plantings will vary but it is important to remember that you are trying to provide connectivity to the landscape adjacent to your property. Don't just look within your property boundaries. If your neighbour's property provides an essential element, such as water, which can be utilized by pollinators visiting your land, you may be able to devote more space to habitat elements that are missing nearby. It is best to use native plants which have evolved to support the needs of specific native pollinators. Some pollinators, however, are generalists and visit many different plants, both native and non-native. Be sure that any non-native plants you choose to use are not invasive. Remember that specialized cultivars sometimes aren't used by pollinators. Flowers that have been drastically altered, such as cultivars with double petals or completely different colour than the wild species, often prevent pollinators from finding and feeding on the flowers. In addition, some cultivars don't contain the same nectar and pollen resources that attract pollinators to the wild types.

• CAUTION: Take time to evaluate the source of your plant material. You want to ensure you get plants that are healthy and correctly identified. Your local garden club can help you make informed decisions when searching for plants.







The following chart lists plants that attract pollinators. It is not exhaustive, but provides guidance on where to start. Annuals, herbs, weeds, and cover crops provide food and shelter for pollinators, too.

Botanical Name	Common Name	Height	Colour	Flower Season	Sun	Soil	Pollinators	Also a host for butterflies
			Trees	and Shrub	S			
Acer saccharum	Sugar maple	12-20m	red, greenish, yellow	March-April	sun to shade	dry	bees, flies	Х
Amorpha fruticosa	False indigo	2.5- 3.5m	purple	May - June	partial shade to shade	wet to moist	bees, flies	
Arctostaphylos uva-ursi	Bearberry	< 1m	white, pink	March - June	sun to shade	dry	butterflies, hummingbirds	Х
Aronia melanocarpa	Black chokeberry	2 m	white	May - June	sun to partial sun	dry to moist	bees, beetles, flies	X
Artemisia campestris	Field sagewort	0.3-1m	green/brown	July - September	sun	dry, well drained	bees	Х
Ceanothus americanus	New Jersey tea	0.5-1m	white	June - August	sun to partial sun	dry	bees, flies, beetles, butterflies	Х
Chamaedaphne calyculata	Leatherleaf	0.3- 1.2m	white/cream	March - June	sun	moist to wet	bees, flies	
Cornus canadensis	Bunchberry dogwood	0.05- 0.2m	white/cream, green/brown	May - September	sun to partial shade	moist, well drained	bees, flies, beetles, butterflies	Х
Cornus stolonifera	Redosier dogwood	1.5-4m	white/cream	May - June	sun to partial shade	moist to wet, well drained	butterflies, bees	Х
Diervilla lonicera	Northern bush- honeysuckle	up to 1m	yellow	June - July	sun to shade	dry to moist	bees, moths	Х
Fagus grandifolia	American beech	0.1-0.3 m	yellow, green/ brown	April - May	partial shade to shade	moist	butterflies	Х
Fraxinus pennsylvanica	Green ash	0.2 m	purple, green/ brown	April - May	sun to partial shade	dry to wet, well drained	butterflies	Х
Hamamelis virginiana	Eastern witch- hazel	3.5- 4.5m	yellow-orange	September - November	sun to shade	moist	moths, beetles	Х
Hudsonia tomentosa	Sand Heather	0.5m	yellow	May - July	sun	well drained	bees, flies, wasps	
Populus balsamifera	Balsam poplar	up to 18m	white, yellow, green	April - June	sun to partial shade	well drained	bees, flies	
Prunus pensylvanica	Pin cherry	7.5-12m	white	May - June	sun	dry to moist, well drained	bees	Х
Prunus serotina	Black cherry	18-27m	white	April - May	shade to sun	moist	bees, flies, wasps	Х
Rhus aromatica	Aromatic sumac	1.2-2m	yellow green	April - May	sun to partial shade	dry to moist, well drained	flies, bees	Х
Ribes triste (Ribes spp.)	Red currant	1-3m	greeenish white, white, pink, red	March - June	sun to shade	moist to dry, well drained	flie, beetles, bees	
Rubus pubescens	Dwarf red blackberry	0.1- 0.4m	white/cream, pink	May - July	sun to shade	dry to wet, well drained	butterflies, bees	
Shepherdia canadensis	Russet buffaloberry	1-3m	yellow, green/ brown	April - May	sun to partial shade	dry to moist, well drained	bees	
Sorbus decora	Showy mountain ash	5-10m	white/cream	June - July	partial shade to shade	moist	bees, butterflies, flies	







Botanical Name	Common Name	Height	Colour	Flower Season			Pollinators	Also a host for butterflies
				Forbs				
Actaea rubra	Red baneberry	<1m	white/cream	May - July	partial shade to shade	moist	bees, flies	
Anaphalis margaritacea	Pearly everlasting	<1m	white/cream	July - September	sun	dry	butterflies, bees	Х
Andromeda glaucophylla	Bog rosemary	<1m	white, pink	May - June	partial shade	wet	bees, flies	
Apocynum androsaemifolium	Spreading dogbane	0.3-2m	white/cream, pink	June - August	sun, partial shade	dry, well drained	butterflies, bees	X
Arenaria stricta	Rock sandwort	0.1- 0.2m	white	May - July	sun to partial shade	dry	bees, flies	
Asclepias incarnata	Swamp milkweed	1.2- 1.5m	pink to reddish	June - October	sun to part shade	moist	flies, butterflies, hummingbirds	Х
Asclepias tuberosa	Butterfly weed	<1m	orange to yellow	May - August	sun	moist to dry	bees, butterflies, flies	Х
Asclepias syriaca	Common milkweed	0.5- 1.5m	purple, pink	June - August	sun	moist, well drained	butterflies	Х
Aster umbellatus (Doellingeria umbellata)	Flat-topped white aster	0.6-2m	white/cream	August - September	sun	moist to wet, well drained	butterflies, bees	Х
Aquilegia canadensis	Red columbine	0.3-1m	red with yellow	April - July	partial shade to sun	moist, well drained	hummingbirds, bees, moths	Х
Caltha palustris	Marsh marigold	0.4- 0.6m	yellow	April - June	sun to partial shade	moist to wet	butterflies	
Campanula rotundifolia	Bluebell bellflower	0.2- 0.5m	blue, purple	June - September	sun to partial shade	dry, well drained	hummingbirds	
Chamerion angustifolium	Fireweed	0.6- 1.8m	purple, pink	July - September	sun	dry to moist, well drained	hummingbirds, butterflies, bees	Х
Cleome serrulata	Rocky mountain beeplant	0.3- 0.8m	white/cream, pink	June - September	sun	dry	bees	
Clintonia borealis	Blue-bead lily	0.1- 0.4m	yellow, green/ brown	May - June	shade	moist	bees	
Coptis groenlandica	Goldthread	0.1- 0.2m	white/cream	May - August	partial shade to shade	moist	bees, flies	
Corydalis sempervirens	Pale corydalis	0.1- 0.6m	yellow, purple, pink	May - September	partial shade	dry	bees, butterflies	
Cypripedium parviflorum var. pubescens	Large yellow ladyslipper	0.5 m	yellow, brown	April - August	shade to sun	moist to dry	bees, flies, beetles	
Epigaea repens	Trailing arbutus	0.1- 0.15m	white/cream, pink	May-June	partial shade	well drained	butterflies	Х
Erigeron annuus	Eastern daisy fleabane	0.3- 1.2m	white/cream	May - October	sun to partial shade	dry, well drained	butterflies, bees	
Eupatorium maculatum	Joe-pye weed	0.6- 1.8m	purple, pink	July-September	sun to partial shade	moist to wet, well drained	butterflies, bees	
Helenium autumnale	Sneezeweed	1-1.5m	yellow, brown	July - September	sun to partial shade	moist to wet	bees, wasps, flies, butterflies	х
Heliopsis helianthoides	False sunflower	1m	yellow	July - October	sun	dry to moderately moist	bees, butterflies	

CONTINUED ON PAGE 18

# PLANTS THAT ATTRACT POLLINATORS IN THE ST. LAWRENCE LOWLANDS

Botanical Name	Common Name	Height	Colour	Flower Season	Sun	Soil	Pollinators	Also a host for butterflies
Impatiens pallida	Jewelweed	up to 2m	white to yellow	June - October	shade	moist	bees	
Iris versicolor	Northern blue flag	0.6- 0.9m	blue, purple	May - August	sun to partial shade	moist to wet	hummingbirds, bees	
Iris virginica var. shrevei	Shreve's iris	0.5-1m	blue, yellow, white	May - July	sun to partial shade	wet to moist	bees	
Kalmia angustifolia	Sheep laurel	0.3- 0.9m	red, white/cream, pink	May - August	sun to partial shade	dry to wet, well drained	butterflies, bees	Х
Kalmia polifolia	Bog laurel	0.1- 0.9m	white/cream, pink	June - August	sun to partial shade	wet	bees	
Ledum groenlandicum	Labrador tea	0.5- 1.5m	white/cream	April - June	sun to shade	moist to wet	butterflies, bees	Χ
Liatris spicata	Blazing star	up to 1.5m	pinkish purple	June - July	sun to partial sun	moist to wet	bees, butterflies, hummingbirds	Χ
Lobelia cardinalis	Cardinal flower	1-1.5m	red	August - October	sun to partial shade	moist to wet	bees, hummingbirds	
Maianthemum canadense	Canada Mayapple	0.05- 0.25m	white/cream	May - June	partial shade to shade	moist, well drained	bees, flies	
Mertensia virginica	Virginia bluebells	up to 0.5m	light blue	May - July	partial sun to shade	moist	bees, butterflies, moths, hummingbirds	Х
Mitella nuda	Naked mitrewort	0.1- 0.2m	white, green	May - August	shade	moist	flies	
Monarda didyma	Beebalm	1-2m	scarlet red	July - September	sun to partial shade	moist to wet	hummingbirds, butterflies, bees	Х
Monarda fistulosa	Wild bergamot	1m	lavendar pink to violet blue	July - September	sun to partial sun	moist	butterflies, hummingbirds, bees, wasps,	Х
Moneses uniflora	One-flowered pyrola	0.1 m	white/cream	June - July	sun to partial shade	dry	bees	
Monotropa uniflora	Indianpipe	0.1-0.2 m	white/cream	June - September	deciduous shade	moist, well drained	bees	
Oligoneuron album	Prairie goldenrod	up to 1m	white	July - September	sun	dry to well drained	bees, flies, butterflies	
Penstemon hirsutus	Hairy beardtongue	<1m	pink to violet blue	June - July	sun to partial sun	moist to dry	bees	Х
Platanthera aquilonis	Northern green orchid	0.05- 0.6m	white, yellow, green	May - August	shade to sun	wet	flies, beetles	
Podophyllum peltatum	Mayapple	<1m	white	March - May	shade	moist to well drained	bees, beetles	
Saxifraga aizoon	White mountain saxifrage	0.1- 0.2m	white/cream, pink	May - June	sun	dry	flies, bees, moths	
Sisyrinchium montanum	Blue-eyed grass	0.1- 0.5m	blue, purple	May - July	sun to shade	dry to wet, well drained	bees	



Botanical Name	Common Name	Height	Colour	Flower Season	Sun	Soil	Pollinators	Also a host for butterflies
Solidago canadensis	Canada goldenrod	0.3- 1.2m	yellow, orange	July - October	sun to partial shade	dry, well drained	butterflies, bees	
Solidago rugosa	Rough- stemmed goldenrod	0.3- 1.8m	yellow	July - October	sun to partial shade	moist to wet	butterflies, bees	
Symphyotrichum novae- angliae	New England aster	up to 1m	purple	September - October	sun to partial shade	moist, well- drained	bees, butterflies, flies	
Symphyotrichum puniceum	Purplestem aster	0.60-2m	blue, purple	August - September	sun	moist to wet, well drained	butterflies, bees	Х
Symplocarpus foetidus	Eastern skunk- cabbage	up to 1m	dark maroon	April - May	sun to shade	wet to flooded	beetles, flies	
Taraxacum ceratophorum	Horned dandelion	0.05- 0.5m	yellow	March - April	sun to partial shade	well drained	bees	
Thalictrum pubescens	Tall meadow- rue	0.6-3m	white/cream	June - August	partial shade	moist	butterflies, bees	
Tiarella cordifolia	Foamflower	<1m	white	April - May	sun to shade	moist	bees, flies, moths	
Zizia aurea	Golden Alexanders	up to 1m	yellow	May - July	sun to partial sun	moist to wet	flies, bees	Х
				Vines				
Celastrus scandens	American bittersweet	2-7m	yellow, green/ brown	May - June	sun to partial shade	dry to moist, well drained	bees	
Clematis virginiana	Virgin's bower	3m+	white	July - August	sun to partial sun	moist	flies, bees	
Parthenocissus quinquefolia	Virginia creeper	15m+	greenish white	May - June	sun to shade	dry to moist	bees, flies	Х
Vitis riparia	Riverbank grape	1-10m	yellow, white/ cream, green/ brown	May - July	sun to shade	dry to moist	bees	
			Sho	elter Plants				
Andropogon gerardii	Big bluestem	1-3.5m	yellow to brown	August - November	sun to partial shade	moist		
Sporobolus cryptandrus	Sand dropseed	0.4- 0.9m	yellow	May - November	sun to partial shade	dry		
Elymus canadensis	Canada wildrye	up to 1m	yellow, green, brown	March - June	sun to partial shade	moist		
Panicum virgatum	Switchgrass	0.4-2m	purple, greenish brown, pink	July - September	sun	dry to moist, well drained	X	
Schizachyrium scoparium	Little bluestem	<1m	yellow, green, brown	June - December	sun to partial shade	dry		
Scirpus cyperinus	Woolgrass	1-1.3m	greenish brown	July - September	sun to partial shade	wet	Х	
Festuca subverticillata	Nodding fescue	0.6m	yellow	April - May	partial shade	dry	Х	

### HABITAT HINTS

### FOR THE ST. LAWRENCE LOWLANDS

BEE-POLLINATED GARDEN FLOWERS AND CROPS											
	Bumble	Digger	Lg Carpenter	Sm Carpenter	Squash/ Gourd	Leafcutter	Mason	Sweat	Plasterer	Yellow- faced	Andrenid
FLOWERS											
Catalpa			Х								
Catnip	х	Х					Х				
Clover		Х									Х
Columbine	х										
Cow parsley										Х	
Goldenrod	х	Х				Х		Х			
Impatiens	Х										
Irises	х		Х								
Lavender	х	Х	х			Х					
Milkwort								Х			
Morning glory				Х							
Penstemon	х	Х					Х				
Phacelia	х	Х		Х		Х	Х	Х	Х		х
Potentilla										Х	
Rose	Х		Х				Х	Х		Х	
Salvia	Х	Х	Х			Х	Х				
Saxifrages								Х		Х	
Sorrel				Х							
Sunflowers	Х	Х	Х	Х		Х		Х	Х		Х
Violet								Х			Х
Wild Mustard		Х							Х		
Willow catkins									Х		Х
					CROPS						
Apple							Х				
Blueberry	х	Х									х
Cherry							Х				х
Eggplant	х		х					Х			
Gooseberry	х										Х
Legumes	х	Х				Х		Х			
Water melon	х							Х			
Squash/ Pumpkins/ Gourds			х		Х						
Tomatoes	х	Х	х					Х			
Thyme	Х	Х					Х	Х		Х	



#### HABITAT AND NESTING REQUIREMENTS:

#### **Honey Bees:**

Honey bees are colonial cavity nesters. Occasionally in the spring and summer you might encounter a swarm of honey bees on the move to a new home. The majority of honey bees are managed by beekeepers in hives. Beekeepers with commercial operations can have bee yards with tens to hundreds of hives commonly in agricultural, rural and some natural habitats. In urban and garden settings it is common to see a single or a handful of honey bee hives — usually wooden boxes painted white or other light colours. Give honey bees space and don't approach their hive. Even beekeepers minimize the amount of time they spend working bees. Honey bees have different feeding needs than native bees. Their colony can last multiple years and they feed on flowers from the beginning of spring bloom to the fall. Honey bees visit crops when they are in bloom and forage on a diversity of wildflowers as well. Honey bees also need plants that produce a large amount of nectar to make honey. Clovers, lavenders, mints, and sages are great nectar sources. Honey bees also like to feed off of the pollen of trees and shrubs such as maples, willows, and roses. Fields of goldenrod are an excellent pollen source.

#### **Bumble Bees:**

Bumble bees nest in cavities such as abandoned mouse nests, other rodent burrows, upside down flower pots, under boards, and other human-made cavities. Colonies are founded by a queen in the spring. The number of workers in a colony can grow to upward of 200 at the peak of summer bloom. Bumble bee colonies die out in the fall after producing new queens. New queens mate and then overwinter, hiding in cracks or small crevices. Bumble bees are usually active during the morning hours and forage at colder temperatures than honey bees, even flying in light rain.

#### Large carpenter bees:

Large carpenter bees chew nests into dead wood, poplar, cottonwood or willow trunks and limbs. The also make nests in structural timbers including redwood and cedar. Depending on the species, there may be one or two brood cycles per year. These bees can be active all day even in the hottest and coolest weather.

#### Digger bees:

Digger bees can be found nesting in sandy, compacted soils, and along stream banksides. These bees are usually active in the morning hours, but can be seen at other times of the day as well. To attract these bees have some areas of exposed soil in your garden and avoid applying thick layers of mulch that are hard to dig through.

#### Small carpenter bees:

Small carpenter bees chew into pithy stems, including roses and blackberry canes, where they make their nests. These bees are more active in the morning but can be found at other times.

#### Squash and Gourd bees:

Squash and gourd bees prefer to nest in sandy soil but also may nest in gardens (where pumpkins, squash and gourds are grown). These bees are early risers and can be found in pumpkin patches before dawn. Males often sleep in wilted squash flowers.

#### **Leaf-cutting bees:**

Leaf-cutting bees nest in pre-existing circular tunnels of various diameters in dead wood created by emerging beetles. Some also nest in the ground. Leafcutter bees line and cap their nests with leaves or flower petals. These bees can be seen foraging throughout the day even in very hot weather.

#### Mason hees

Mason bees use pre-existing tunnels of various diameters in dead wood made by emerging beetles, or human-made nesting substrates such as drilled wood blocks or cardboard tubes. These bees get their name from the fact that they cap their nests with mud. Mason bees are generally more active in the morning hours.

#### Sweat bees:

Sweat bees need bare ground in sunny areas not covered by vegetation for nesting. Some will nest in small pre-existing holes, much like leaf-cutting or mason bees. Like most bees, sweat bees forage for pollen earlier in the morning and then for nectar later in the day. To help these bees nest, keep some parts of your garden exposed and avoid applying thick layers of mulch that are hard to dig through.

#### Plasterer or cellophane bees:

Plasterer or cellophane bees get their name from the unique, clear waterproof lining they make around their nest. Similar to sweat bees they prefer bare ground, stream banks or slopes. These bees can be active in the morning or later in the day.

#### Yellow-faced bees:

Yellow-faced bees are tiny, hairless bees that make their nesting by chewing into small dead stems. These bees are more active during morning hours.

#### Mining bees:

Mining bees prefer sunny, bare ground, and sand soil. They are also known to nest under leaf litter or in the soil along banksides and cliffs. Mining bees are active in the spring and most commonly seen on flowers during the morning when pollen and nectar resources are abundant.

### A BASIC CHECKLIST

# S.H.A.R.E. - SIMPLY HAVE AREAS RESERVED FOR THE ENVIRONMENT.

- Planting for pollinators is the first step. Put your pollinator habitat on the S.H.A.R.E. map and connect to pollinator conservation efforts across North America. Visit pollinator.org/SHARE
- Farmers can also certify their habitat with Bee-Friendly Farming. Visit pollinator.org/bff

# BECOME FAMILIAR WITH POLLINATORS IN YOUR LANDSCAPE.

- Watch for activity throughout the day and the seasons.
- Keep a simple notebook of when and what comes to your garden.
- **%** Take part in citizen science monitoring and get involved in local programs.
- Consult a local field guide or web site when you are ready to learn more details.

# ADD NATIVE PLANTS TO ATTRACT MORE NATIVE POLLINATORS.

- **%** List the plants you currently have in your landscape.
- Determine when you need additional flowers to provide nectar and pollen throughout the growing season.
- Add plants that provide additional seasons of bloom, create variable heights for shelter, and attract the types of pollinators you want.
- Mon't forget to include host plants that provide food and shelter for larval development.
- **%** Contact your local native plant society or extension agent for more help.

# USE POLLINATOR FRIENDLY LANDSCAPE PRACTICES TO SUPPORT THE POLLINATORS YOU ATTRACT.

- We Integrated Pest Management (IPM) practices to address pest concerns.
- Tolerate a little mess leave dead snags and leaf litter, keep areas bare for ground nesting insects, and leave some weeds that provide food for pollinators.
- Provide safe access to clean water.

# NOTICE THE CHANGES THAT YOU HAVE HELPED TO CREATE!





Many books, websites, and people were consulted to gather information for this guide. Use this list as a starting point to learn more about pollinators and plants in your area.

#### **ECOREGION PROFILE**

#### St. Lawrence Lowlands

http://www.pc.gc.ca/eng/docs/v-g/nation/nation76.aspx

Environment Canada Ecozones Program - St. Lawrence Lowlands http://ecozones.ca/english/region/132

#### Government of Ontario - Species at Risk

https://www.ontario.ca/page/species-risk

#### POLLINATION/POLLINATORS

#### Pollinator Partnership

www.pollinator.org

#### North American Pollinator Protection Campaign

www.nappc.org

#### Pollination Canada

www.pollinationcanada.ca

#### Seeds of Diversity

www.seeds.ca

### Canadian Biodiversity Information Facility: Butterflies of Canada

www.cbif.gc.ca/eng/species-bank/butterflies-of-canada/?id=1370403265518

#### North American Butterfly Association

www.naba.org

#### Canadian Honey Council

www.honeycouncil.ca

# Buchmann, S.L. and G.P. Nabhan. 1997. The Forgotten Pollinators Island Press: Washington, DC.

# Committee on the Status of Pollinators in North America. 2007. Status of Pollinators in North America The National Academies Press: Washington, DC.

#### **NATIVE PLANTS**

#### Montreal Botanical Garden

http://espacepourlavie.ca

#### **Ontario Wildflowers**

www.ontariowildflowers.com

#### Fletcher Wildlife Garden

www.ofnc.ca/fletcher/

#### Roger Van den Hende Garden

http://www.jardin.ulaval.ca/en/

### **FEEDBACK**

We need your help to create better guides for other parts of North America. Please e-mail your input to feedback@pollinator.org.

- **%** How will you use this guide?
- **%** Do you find the directions clear? If not, please tell us what is unclear.
- Is there any information you feel is missing from the guide?
- **%** Any other comments?

THANK YOU
FOR TAKING
THE TIME TO HELP!



**RESEARCH AND WRITING:** 

VICTORIA WOJCIK

AMBER BARNES

EDITORIAL:

AMBER BARNES, VICTORIA WOJCIK

DESIGN:

**MARGUERITE MEYER** 

FUNDING:

Stanley Smith Horticultural Trust

PHOTO CONTRIBUTORS:

Lara Roketenetz, Tim Dunne, Balalaika

ILLUSTRATIONS:

Carolyn Vibbert